

This Page Is Inserted by IFW Operations  
and is not a part of the Official Record

## BEST AVAILABLE IMAGES

Defective images within this document are accurate representations of  
the original documents submitted by the applicant.

Defects in the images may include (but are not limited to):

- BLACK BORDERS
- TEXT CUT OFF AT TOP, BOTTOM OR SIDES
- FADED TEXT
- ILLEGIBLE TEXT
- SKEWED/SLANTED IMAGES
- COLORED PHOTOS
- BLACK OR VERY BLACK AND WHITE DARK PHOTOS
- GRAY SCALE DOCUMENTS

## IMAGES ARE BEST AVAILABLE COPY.

As rescanning documents *will not* correct images,  
please do not report the images to the  
Image Problem Mailbox.

PCT

WORLD INTELLECTUAL PRO  
International

WO 9606077A1

INTERNATIONAL APPLICATION PUBLISHED UNDER



CS

(51) International Patent Classification 6 : C07D 207/12, 207/32, 401/12, 405/12, 409/12, A61K 31/40  
A1  
(11) International Publication Number: WO 96/06077  
(43) International Publication Date: 29 February 1996 (29.02.96)

(21) International Application Number: PCT/IB95/00374  
(22) International Filing Date: 18 May 1995 (18.05.95)  
(30) Priority Data: PCT/JP94/01399 24 August 1994 (24.08.94) WO  
(34) Country for which the regional or international application was filed: JP

(71) Applicant (for JP only): PFIZER PHARMACEUTICALS INC.  
[JP/JP]; Shinjuku Mitsui Building, 1-1, Nishi-shinjuku 2-chome, Shinjuku-ku, Tokyo 163-04 (JP).

(71) Applicant (for all designated States except JP US): PFIZER INC. [US/US]; 235 East 42nd Street, New York, NY 10017 (US).

(72) Inventor; and (75) Inventor/Applicant (for US only): ITO, Fumitaka [JP/JP]; 1-2, Sakuragawa 1-chome, Takekoyo-cho, Chita-gun, Aichi 470-23 (JP).

(74) Agents: SPIEGEL, Allen, J. et al.; Pfizer Inc., Patent Dept., 235 East 42nd Street, New York, NY 10017 (US).

(54) Title: N-[2-(PYRROLIDINYL-1)-1-PHENYLETHYL]ACETAMIDES AS KAPPA RECEPTOR ANTAGONISTS

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

(54) Title: N-[2-(PYRROLIDINYL-1)-1-PHENYLETHYL]ACETAMIDES AS KAPPA RECEPTOR ANTAGONISTS

(54) Title: N-[2-(PYRROLIDINYL-1)-1-PHENYLETHYL]ACETAMIDES AS KAPPA RECEPTOR ANTAGONISTS

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

(57) Abstract  
A compound of formula (I) and its pharmaceutically acceptable salt, wherein R is hydrogen or hydroxy; Ar is unsubstituted or substituted phenyl; X is unsubstituted or substituted phenyl or heterocyclic, mono-, di- or trihalomethyl, cyano, or the like; and X' is phenyl, naphthyl, furyl, thieryl, pyridyl, thiazolyl, benzofuryl, benzothienyl or the like. These compounds have agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic, and neuroprotective agent.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AT	Austria	GB	United Kingdom	MR	Mauritania
AU	Australia	GE	Georgia	MW	Malawi
BB	Barbados	GN	Guinea	NE	Niger
BE	Belgium	GR	Greece	NL	Netherlands
BF	Burkina Faso	HU	Hungary	NO	Norway
BG	Bulgaria	IE	Ireland	NZ	New Zealand
BJ	Benin	IT	Italy	PL	Poland
BR	Brazil	JP	Japan	PT	Portugal
BY	Belarus	KE	Kenya	RO	Romania
CA	Canada	KG	Kyrgyzstan	RU	Russian Federation
CF	Central African Republic	KP	Democratic People's Republic of Korea	SD	Sudan
CG	Congo	KR	Republic of Korea	SE	Sweden
CH	Switzerland	KZ	Kazakhstan	SI	Slovenia
CI	Côte d'Ivoire	LJ	Liechtenstein	SK	Slovakia
CM	Cameroon	LK	Sri Lanka	SN	Senegal
CN	China	LU	Luxembourg	TD	Chad
CS	Czechoslovakia	LV	Latvia	TG	Togo
CZ	Czech Republic	MC	Monaco	TJ	Tajikistan
DE	Germany	MD	Republic of Moldova	TT	Trinidad and Tobago
DK	Denmark	MG	Madagascar	UA	Ukraine
ES	Spain	ML	Mali	US	United States of America
FI	Finland	MN	Mongolia	UZ	Uzbekistan
FR	France			VN	Viet Nam
GA	Gabon				

-1-

N-(2-(pyrrolidinyl-1)-1-phenylethyl) acetamides as Kappa receptor antagonists

### Technical Field

This invention relates to novel carboxamide compounds and their pharmaceutically acceptable salts, and to pharmaceutical compositions containing them. These compounds and compositions are useful as analgesic, antiinflammatory, diuretic or  
10 neuroprotective agents for the treatment of a mammalian subject, especially a human subject.


### Background Art

Opioid analgesics such as morphine are therapeutically useful, but their usage is strictly limited because of their side effects such as drug dependency. Thus, analgesics  
15 with high usefulness and reduced tendency to cause drug dependency are desired. Considerable pharmacological and biochemical studies have been carried out to discover the opioid peptides and opioid receptors, and the discovery of the subtype of opioid receptor such as mu, delta, kappa at a peripheral nerve in a variety of species, including human, has made a beginning towards creating new analgesics. As it is  
20 thought that opioid analgesics such as morphine act as a  $\mu$ -receptor agonist, separating the action based on a kappa-receptor agonist from the action based on  $\mu$ -receptor agonist has been investigated. Recently kappa-selective agonists have been reported from the above viewpoint for example, EMD-60400: A. Barber et al., Naunyn-Schmied. Arch. Pharmacol., 345 (Suppl.): Abst 456. Some of them actually have been  
25 studied in clinical trials (Med. Res. Rev., 12, 525 (1992)).

However, even when a selective kappa-receptor agonist is employed, use of high doses can give rise to side effects such as sedation. Therefore, it would be desired to provide compounds having better agonist activity toward opioid kappa receptor.

### Brief Disclosure of the Invention

30 The present invention provides a compound of the following formula:



-2-


and its pharmaceutically acceptable salt, wherein

R is hydrogen or hydroxy;

Ar is phenyl or phenyl substituted with one to three substituents selected from halo, C<sub>1-4</sub> alkyl and C<sub>1-4</sub> alkoxy;

- 5 X is phenyl or heterocyclic; phenyl or heterocyclic substituted with one to three substituents selected from halo, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy and methoxycarbonyl; mono-, di- or tri-halomethyl; cyano; COR<sup>1</sup>, CH=NOR<sup>2</sup>, OR<sup>2</sup>, SR<sup>2</sup>, CH<sub>2</sub>CN, CH<sub>2</sub>OR<sup>2</sup>, CH<sub>2</sub>SR<sup>2</sup>, CH<sub>2</sub>S(O)R<sup>2</sup>, CH<sub>2</sub>S(O)<sub>2</sub>R<sup>2</sup>, CH<sub>2</sub>(R<sup>2</sup>)R<sup>3</sup>, CH<sub>2</sub>N(R<sup>2</sup>)R<sup>3</sup>, CH<sub>2</sub>NR<sup>2</sup>OH, CH<sub>2</sub>N(COR<sup>2</sup>)OH, CH<sub>2</sub>NR<sup>2</sup>COR<sup>3</sup>, CH<sub>2</sub>NR<sup>2</sup>S(O)<sub>2</sub>R<sup>3</sup> or CH<sub>2</sub>OCOR<sup>2</sup>,  
 10 wherein R<sup>1</sup> is hydrogen, hydroxy, amino, NHOH, NHOCH<sub>3</sub>, pyridylamino, NHN(CH<sub>3</sub>)<sub>2</sub>, C<sub>1-4</sub> alkoxy, benzyloxy, C<sub>1-4</sub> alkylamino, di-C<sub>1-4</sub> alkylamino, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkylthio; and R<sup>2</sup> and R<sup>3</sup> are each hydrogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy or C<sub>7-11</sub> phenylalkyl; and  
 X<sup>1</sup> is phenyl, naphthyl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl or benzothienyl;  
 15 phenyl, naphthyl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl or benzothienyl, substituted with one to three substituents selected from halo, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, amino, hydroxy, nitro, trifluoromethyl and mesyl.

Further, the present invention provides a compound of the formula:



- 20 wherein R, Ar and X are as already defined. These compounds can be used as intermediates to prepare the compounds of formula (I).

The carboxamide compounds of the present invention of formula (I) exhibit significant agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic and neuroprotective agents, in mammals, especially man.

- 25 Accordingly, the present invention also provides a pharmaceutical composition useful as an analgesic, antiinflammatory, diuretic or neuroprotective agent, in a mammal, especially man, which comprises a therapeutically effective amount of the carboxamide

compound of formula (I) or its pharmaceutically acceptable salt together with a pharmaceutically acceptable carrier.

#### Detailed Disclosure of the Invention

In this specification, the term "heterocyclic" means a monocyclic or bicyclic hydrocarbon group which has one or more hetero atoms in the ring, preferably has 4 to 10 carbon atoms and 1 to 3 heteroatoms, including piperidino, morpholino, thiamorpholino, pyrrolidino, pyrazolino, pyrazolidino, pyrazoryl, piperaziny, furyl, benzofuryl, thienyl, benzothienyl, oxazolyl, tetrazolyl, thiazolyl, imidazolyl, pyrazolyl, pyridyl, pyrimidinyl, pyrrolyl, pyrrolidinyl, quinolyl and quinuclidinyl.

A preferred group of compounds of this invention includes the compounds of formula (I) wherein R is hydroxy; Ar is phenyl optionally substituted with one to three halogen atoms, preferably phenyl; X is phenyl optionally substituted with one to three substituents selected from halo, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy and methoxycarbonyl; and X<sup>1</sup> is phenyl optionally substituted with one to three halogen atoms, preferably 3,4-dichlorophenyl.

Another preferred group of compounds of this invention includes the compounds of formula (I) wherein R is hydroxy; Ar is phenyl optionally substituted with one to three halogen atoms, more preferably phenyl; X is mono-, di- or tri-halomethyl, cyano, hydroxycarbonyl, butyloxycarbonyl, benzyloxycarbonyl, carbamoyl or hydroxymethyl; and X<sup>1</sup> is phenyl optionally substituted with one to three halogen atoms, preferably 3,4-dichlorophenyl.

Another preferred group of compounds of this invention includes the compounds of formula (I) wherein R is hydroxy; Ar is phenyl optionally substituted with one to three halogen atoms, more preferably phenyl; X is furyl, thienyl, pyridyl or oxadiazolyl; and X<sup>1</sup> is phenyl optionally substituted with one to three halogen atoms, preferably 3,4-dichlorophenyl.

Preferred individual compounds of the invention are:

- N-carboxymethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;
- 2-(3,4-dichlorophenyl)-N-(2-hydroxyethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;
- 2-(3,4-dichlorophenyl)-N-[(2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl)-N-(2,2,2-trifluoroethyl)acetamide];

2-(3,4-dichlorophenyl)-N-furfuryl-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

2-(3,4-dichlorophenyl)-N-[2-(3-(s)-hydroxypyrrolidin-1-yl)-1-(s)-phenylethyl]-N-(4-pyridyl)methylacetamide;

5 2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(3-pyridyl)methylacetamide;

N-cyanomethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

2-(3,4-dichlorophenyl)-N-(2,2-difluoroethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;

10 N-2-cyanoethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide;


2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-methoxycarbonylmethylacetamide; and

15 2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(1,2,4-oxadiazol-3-yl)methylacetamide.

#### General Synthesis

The carboxamide compounds of formula (I) of this invention may be prepared by a variety of synthetic methods. For example, the carboxamide compounds of formula (I) may be prepared by acylation of compound (II), as indicated in the following Preparation Method A-I.

#### Preparation Method A-I:



(wherein R, Ar, X and X' are as previously defined)

25 In Preparation Method A-I, the amine compound (II) is reacted with an acylating agent using standard acylating techniques known to those skilled in the art. In a typical acylation method, the amine compound (II) may be reacted with acyl halide (e.g., X'CH<sub>2</sub>COCl) in a suitable reaction-inert solvent. Suitable inert-reaction solvents include,


-5-

for example, aromatic hydrocarbons such as benzene, toluene and xylene; ethers such as ethyl ether, dioxane and tetrahydrofuran; halogenated hydrocarbons such as chloroform, dichloromethane and dichloroethane; amides such as N,N-dimethylformamide; and nitriles such as acetonitrile. If desired, this reaction may be catalyzed by a base such as triethylamine, pyridine or alkoxide. The reaction may be carried out at a temperature of from -30°C to 100°C, preferably from 0°C to 25°C, for 10 minutes to 48 hours, preferably from 30 minutes to 24 hours.

The compound (I) of the present invention may also be obtained from the amine compound (II) by the other acylation methods, for example, (1) a reaction with anhydride (e.g.,  $(X^1CH_2CO)_2O$ ) or a mixed anhydride in the presence of base; (2) a reaction with carboxylic acid ( $X^1CH_2COOH$ ) in the presence of a coupling agent such as dicyclohexylcarbodiimide (DCC), water soluble carbodiimide (WSCD), 2-ethoxy-N-ethoxycarbonyl-1,2-dihydroquinoline, Bop agent (Benzotriazol-1-yloxy-tris(dimethylamino)phosphonium hexafluorophosphate), diethyl azodicarboxylate-triphenylphosphine, diethyl cyanophosphonate, carbonyldiimidazole and diphenylphosphoryl azide; or (3) a reaction with carboxylic ester (e.g.,  $X^1CH_2COOR'$  wherein  $R'$  is lower alkyl) optionally in the presence of base. The conditions employed for the acylation methods can be properly chosen by the skilled persons.

In an alternative method, the compound (I) of the present invention may be prepared by the following Preparation Method A-II.

Preparation Method A-II:



In this method, the compound (I) may be obtained by alkylation of the amide compound (III). Alkylation methods known to those skilled in the art can be used. For example, the amide compound (III) may be reacted with alkylhalide (e.g.,  $XCH_2L$  wherein  $X$  is as previously defined; and  $L$  is halogen such as chlorine) in a reaction-inert solvent. If desired, this reaction may be catalyzed by a base such as sodium, sodium hydride, sodium hydroxide, potassium hydroxide, with or without a phase-transfer




-6-

catalyst. The reaction may be carried out at a temperature of from 0°C to 200°C, preferably from 60°C to 150°C, for 5 minutes to 24 hours, preferably from 30 minutes to 12 hours.

Alternatively, the alkylation of the compound (III) may be carried out by reacting the compound (III) with formaldehyde and metal salts (e.g., MX wherein X is as previously defined; and M is an alkali metal such as sodium and potassium) in a suitable reaction-inert solvent. In addition, the amide compound (III) may be obtained by acylation of the amine compound (IV) in similar procedures to those described in Preparation Method A-I above.

10 In the present invention, the amine compound (II) may be obtained by the following Preparation Method B-I.

Preparation Method B-I:




In this method, the amine compound (II) may be obtained by alkylation of the amine compound (IV) using standard alkylation techniques known to those skilled in the art. A preferred alkylation method is reductive alkylation wherein the amine compound (IV) may be reacted with aldehyde, XCHO (wherein X is as already defined) in the presence of a reducing agent such as NaBH<sub>4</sub>, NaBH<sub>3</sub>CN or NaBH(OAc)<sub>3</sub>. This reaction may be carried out in a suitable reaction-inert solvent at a temperature of from -20°C to 60°C, preferably from 0°C to 25°C, for 10 minutes to 48 hours, preferably from 60 minutes to 5 hours. In an alternative alkylation method, the amine compound (II) may be obtained by reacting the amine compound (IV) with alkylhalide, XCH<sub>2</sub>L (wherein X and L are as already defined) under conditions known to those skilled in the art. The Mannich type alkylation can be also used, which comprises the reaction of the compound (IV) with formaldehyde and a metal salt. The amine compounds (IV) are either known or may be prepared by known methods as described in European Patent No. 254545.

Alternatively, the amine compound (II) may be obtained by acylation of the compound (IV), followed by reduction, as indicated in the following Preparation Method B-II.

Preparation Method B-II:


-7-



In a typical procedure, the amine compound (IV) may be first reacted with acylating agents,  $\text{XCOOH}$  (wherein X is as already defined) in the presence of a suitable coupling agent as mentioned above, in a suitable reaction-inert solvent, followed by reduction using a reducing agent such as  $\text{LiAlH}_4$ ,  $\text{BH}_3 \cdot \text{Me}_2\text{S}$  or  $\text{BH}_3 \cdot \text{THF}$ . This reaction may be carried out at a temperature of from  $0^\circ\text{C}$  to  $100^\circ\text{C}$ , preferably from  $20^\circ\text{C}$  to  $80^\circ\text{C}$ , for 30 minutes to 24 hours, preferably from 60 minutes to 12 hours. The other possible acylation methods prior to the reduction include a reaction of the compound (IV) with acyl halide,  $\text{XCOL}$  in the presence of base; and the reaction of the compound (IV) with anhydride,  $(\text{XCO})_2\text{O}$  in the presence of base. The conditions to be employed for these acylation methods can be appropriately chosen by those skilled in the art.

Further, the compound (II) may be obtained by acylation of an amide compound of the following formula (V), followed by reduction, as indicated in the following Preparation Method B-III.

15 Preparation Method B-III:



In this method, the amide compound (V) may be first subjected to acylation as mentioned in the above Preparation Method B-II, and then subjected to reduction, to obtain the compound (II). The conditions for this reaction may be similar to those described in the above Preparation Method B-II. In addition, the amide compound (V) is either known or can be prepared by known methods as described in, for example, European Patent N . 254545 and Chem. Pharm. Bull., 42(3) 690-693, 1994.

The compounds of formula (I), and the intermediates shown in the above Preparation Methods can be isolated and purified by conventional procedures, such as recrystallisation or chromatographic purification.

As the carboxamide compounds of this invention possess at least two asymmetric centers, they are capable of occurring in various stereoisomeric forms or configurations. Hence, the compounds can exist in separated (+)- and (-)-optically active forms, as well as mixtures thereof. The present invention includes all such forms within its scope. Individual isomers can be obtained by known methods, such as optically selective reaction or chromatographic separation in the preparation of the final product or its intermediate.

The carboxamide compounds of the present invention can be used in the form of the inorganic salts with acid such as hydrochloric acid, hydrobromic acid, sulfonic acid, nitric acid, phosphoric acid and the like and the organic salts with acid such as acetic acid, formic acid, benzoic acid, oxalic acid, succinic acid, fumaric acid, maleic acid, citric acid, alkylsulfonic acid.

The carboxamide compounds of the present invention of formula (I) exhibit significant agonist activity toward opioid kappa receptor and are thus useful as analgesic, antiinflammatory, diuretic and neuroprotective agents for the treatment of mammals, especially humans in need of such agents.

The activity of the carboxamide compounds of formula (I) of the present invention as opioid kappa agonist, is demonstrated by the opioid receptor binding activity. Such activity may be determined in homogenates from guinea pig whole brain, as described by Regina, A. et al. in J. Receptor Res. 12: 171-180, 1992. In summary, tissue homogenate is incubated at 25°C for 30 min in the presence of labelled ligand and test compounds. The  $\mu$ -sites are labelled by 1 nM of (3H)-[D-Ala<sup>2</sup>,MePhe<sup>4</sup>,Gly-ol<sup>5</sup>]enkephalin (DAMGO), the  $\delta$ -sites by 1 nM of (3H)-[D-Pen<sup>2</sup>,5]enkephalin (DPDPE) and the  $\kappa$ -sites by 0.5 nM (3H)-CI-977. The non specific binding is measured by use of 1 mM CI-977 ( $\kappa$ ), 1 mM (DAMGO) ( $\mu$ ), 1mM (DPDPE) ( $\delta$ ). Data are expressed as the IC<sub>50</sub> values obtained by a non-linear fitting program using the Cheng and Prusoff equation. All compounds prepared in the Working Examples as described below were tested by this method, and showed an IC<sub>50</sub> value of 0.01 nM to 10 $\mu$ M with respect to inhibition of binding at its receptor.

The agonist activity toward opioid kappa receptor can also be demonstrated by the Formalin Test as described by Wheeler-Aceto, H. et al. in Psychopharmacology 104: 35-44, 1991. In this

testing, male SD rats (80-100 g) are injected s.c. with a test compound dissolved in 0.1% methyl cellulose saline or vehicle. After 30 min., 50  $\mu$ l of a 2% formalin are injected into a hind paw. The number of licking the injected paw per observation period is measured 15-30 min. after the injection of formalin and expressed as % inhibition compared to the respective vehicle group.

- 5 The agonist activity toward opioid kappa receptor can also be demonstrated by the Rotarod Test as described by Hayes, A.G. et al. in Br. J. Pharmacol. 79: 731-736, 1983. In this testing, a group of 6-10 male SD rats (100-120 g) are selected for their ability to balance on a rotating rod (diameter 9 cm, rate of rotation 5 r.p.m.). The selected rats are then injected s.c. with a test compound dissolved in 0.1% methyl cellulose saline. The animals are tested again 30 min. after treatment; a rat falling off the bar more than twice within 150 seconds is considered to be showing motor impairment and the animal's performance (i.e., time on the rotarod) are recorded. The ED<sub>50</sub> value, defined as the dose of the drug which halves the performance time is observed in the control group.

- 15 The carboxamide compounds of formula (I) of this invention can be administered via either the oral, parenteral or topical routes to mammals. In general, these compounds are most desirably administered to humans in doses ranging from 0.01 mg to 100 mg per day, although variations will necessarily occur depending upon the weight and condition of the subject being treated, the disease state being treated and the particular route of administration chosen. However, a dosage level that is in the range of from 0.01 mg to 50 mg per kg of body weight per day is most desirably employed for the treatment of pain in a postoperative patient.

- 25 The compounds of the present invention may be administered alone or in combination with pharmaceutically acceptable carriers or diluents by either of the above routes previously indicated, and such administration can be carried out in single or multiple doses. More particularly, the novel therapeutic agents of the invention can be administered in a wide variety of different dosage forms, i.e., they may be combined with various pharmaceutically acceptable inert carriers in the form of tablets, capsules, lozenges, troches, hard candies, powders, sprays, creams, salves, suppositories, jellies, gels, pastes, lotions, ointments, aqueous suspensions, injectable solutions, elixirs, syrups, and the like. Such carriers include solid diluents or fillers, sterile aqueous media and various nontoxic organic solvents, etc. Moreover, oral pharmaceutical compositions can be suitably sweetened and/or flavored. In general, the therapeutically-effective compounds of this invention are present in such dosage forms at concentration levels ranging 5% to 70% by weight, preferably 10% to 50% by weight.

For oral administration, tablets containing various excipients such as microcrystalline cellulose, sodium citrate, calcium carbonate, dipotassium phosphate and glycine may be employed along

-10-

with various disintegrants such as starch and preferably corn, potato or tapioca starch, alginic acid and certain complex silicates, together with granulation binders like polyvinylpyrrolidone, sucrose, gelatin and acacia. Additionally, lubricating agents such as magnesium stearate, sodium lauryl sulfate and talc are often very useful for tableting purposes. Solid compositions of a similar type may also be employed as fillers in gelatine capsules; preferred materials in this connection also include lactose or milk sugar as well as high molecular weight polyethylene glycols. When aqueous suspensions and/or elixirs are desired for oral administration, the active ingredient may be combined with various sweetening or flavoring agents, coloring matter or dyes, and, if so desired, emulsifying and/or suspending agents as well, together with such diluents as water, ethanol, propylene glycol, glycerin and various like combinations thereof.

For parenteral administration, solutions of a compound of the present invention in either sesame or peanut oil or in aqueous propylene glycol may be employed. The aqueous solutions should be suitably buffered (preferably pH > 8) if necessary and the liquid diluent first rendered isotonic. These aqueous solutions are suitable for intravenous injection purposes. The oily solutions are suitable for intra-articular, intra-muscular and subcutaneous injection purposes. The preparation of all these solutions under sterile conditions is readily accomplished by standard pharmaceutical techniques well-known to those skilled in the art. Additionally, it is also possible to administer the compounds of the present invention topically when treating inflammatory conditions of the skin and this may preferably be done by way of creams, jellies, gels, pastes, ointments and the like, in accordance with standard pharmaceutical practice.

### Examples

The present invention is illustrated by the following examples. However, it should be understood that the invention is not limited to the specific details of these examples. Melting points were taken with a Buchi micro melting point apparatus and uncorrected. Infrared Ray absorption spectra (IR) were measured by a Shimadzu infrared spectrometer (IR-470). <sup>1</sup>H and <sup>13</sup>C nuclear magnetic resonance spectra (NMR) were measured in CDCl<sub>3</sub> by a JEOL NMR spectrometer (JNM-GX270, 270MHz) unless otherwise indicated and peak positions are expressed in parts per million (ppm) downfield from tetramethylsilane. The peak shapes are denoted as follows: s, singlet; d, doublet; t, triplet; m, multiplet; br, broad.

### Preparation 1

#### (S)-Phenylglycyl-3-(S)-hydroxypyrrolidine

To a stirred solution of 3-(S)-pyrrolidinol (3.054g, 35mmol) and N-benzoyloxycarbonyl-(S)-phenylglycine (10.00g, 35mmol) in DMF (40ml) was added diethyl phosphorocyanidate (6.28ml, 42mmol) followed by addition of N-methylmorpholine (4.65ml, 42mmol) at room temperature.

After 1h stirring at room temperature, the reaction mixture was poured into water (200ml), extracted with mixed solvent (ethyl acetate / hexane / ether : 2/1/1, 100ml x 3). The extract combined was washed with 1N HCl solution, saturated NaHCO<sub>3</sub> aqueous solution and brine, dried (MgSO<sub>4</sub>), and concentrated to give 11.395g (91.9%) of N-benzyloxycarbonyl-(S)-phenylglycyl-3-(S)-hydroxypyrrolidine as yellow oil. A suspension mixture of this oil (11.395g, 32mmol) and 10% palladium carbon (1.14g) in methanol (100ml) was stirred under hydrogen atmosphere at room temperature for 17h. The catalyst was removed by Celite filtration and the filtrate was concentrated to give 8.255g (crude 100%) of the title compound as brown oil.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.45-7.27 (5H, m), 4.56 (0.7H, s), 4.49 (0.3H, s), 4.45-4.35 (1H, m), 3.74-3.36 (3H, m), 3.25-3.15 (0.7H, m), 3.10-2.98 (0.3H, m), 2.70 (2H, br.s), 2.30 (1H, br.s), 2.05-1.75 (2H, m).

IR(neat) : 3350, 3300, 1640cm<sup>-1</sup>.

To a stirred solution of 3-(S)-pyrrolidinol (14.70g, 169mmol) and N-t-butoxycarbonyl-(S)-phenylglycine(42.46g, 169mmol) in DMF (250ml) was added diethyl phosphorocyanidate (30.3ml, 203mmol) followed by addition of N-methylmorpholine (22.5ml, 203mmol) at room temperature. After 1h stirring at room temperature, the reaction mixture was poured into water (1250ml), extracted with mixed solvent (ethyl acetate/hexane/ether:2/1/1, 400ml x 3). The extract combined was washed with 1N HCl solution, saturated NaHCO<sub>3</sub> aqueous solution and brine, dried (MgSO<sub>4</sub>), and concentrated to give 34.81g (64.3%) of N-t-butoxycarbonyl-(S)-phenylglycyl-3-(S)-hydroxypyrrolidine as white powder. To a stirred suspension of N-t-butoxycarbonyl-(S)-phenylglycyl-3-(S)-hydroxypyrrolidine (10.13g, 31.6mmol) in CH<sub>2</sub>Cl<sub>2</sub> (10ml) was added trifluoroacetic acid (25ml) at 0°C and resulting solution was stirred at room temperature for 1h. After evaporation of excess trifluoroacetic acid and solvent, the residue was basified with aqueous NH<sub>3</sub> solution (20ml) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (30ml x 3). After dry (Na<sub>2</sub>SO<sub>4</sub>), the solvent was evaporated to give 4.974g (71.5%) of title compound.

#### Preparation 2

##### (2S,3S)-1-(2-Amino-2-phenylethyl)-3-hydroxypyrrolidine

To a stirred suspension of lithium aluminum hydride (3.795g, 100mmol) in THF (200 ml) was added a solution of (S)-phenylglycyl-3-(S)-hydroxypyrrolidine (7.049g, 32mmol) in THF (100ml) dropwise at room temperature. The reaction mixture was refluxed for 1.5h. Then the reaction mixture was cooled down to room temperature and Na<sub>2</sub>SO<sub>4</sub>·10H<sub>2</sub>O (10.31g) and KF (1.86g) was added to the reaction mixture. After 1h stirring, the white solid precipitated was removed by Celite filtration and the filtrate was concentrated to give 4.21g of clear yellow oil. This was

-12-

purified by column chromatography (silica gel:180g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH/NH<sub>4</sub>OH : 50/5/1 as eluent) to afford 3.584g (54.3%) of clear yellow oil.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.38-7.15 (5H, m), 4.37-4.29 (1H, m), 4.08 (1H, dd, J = 4.0, 10.3Hz), 3.08-3.01 (1H, m), 2.77 (1H, dd, J = 10.3, 12.1 Hz), 2.69-2.61 (1H, m), 2.43 (1H, dd, J = 4.0, 12.1Hz), 2.31-2.00 (6H, m), 1.83-1.70 (1H, m).

IR (neat) : 3350, 3200cm<sup>-1</sup>.

### Preparation 3

#### (2S,3S)-1-[2-N-(Benzyloxycarbonyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine

A mixture of (2S,3S)-1-(2-amino-2-phenylethyl)-3-hydroxypyrrolidine (1.65g, 8mmol), benzyl 2-bromoacetate (1.52ml, 9.6mmol), and triethylamine (1.34ml, 9.6mmol) in CH<sub>2</sub>Cl<sub>2</sub> (30ml) was refluxed for 7h. The reaction mixture was diluted with water (50ml), then extracted with CH<sub>2</sub>Cl<sub>2</sub> (50ml x 3). The extract combined was washed with brine and dried (MgSO<sub>4</sub>). Evaporation of the solvent gave 2.18g of brown oil which was purified by column chromatography (silica gel:70g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH : 20/1 to 10/1 as eluent) to afford 1.218g (42.9%) of clear yellow oil.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.40-7.22 (10H, m), 5.13 (2H, s), 4.35-4.25 (1H, m), 3.82 (1H, dd, J = 3.7, 11.0Hz), 3.43 (1H, d, J = 17.6Hz), 3.22 (1H, d, J = 17.6Hz), 3.16-3.06 (1H, m), 2.86 (1H, dd, J = 11.0, 12.1Hz), 2.70 (1H, dd, J = 4.8, 10.3Hz), 2.64 (1H, dd, J = 1.8, 9.9Hz), 2.31 (1H, dd, J = 3.7, 12.1Hz), 2.26-2.15 (4H, m), 1.82-1.71 (1H, m).

IR (neat) : 3350, 1740cm<sup>-1</sup>.

### Example 1

#### N-(Benzyloxycarbonyl)methyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

To a stirred solution of (2S,3S)-1-[2-N-(benzyloxycarbonyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine (354g, 1mmol) in dioxane (4ml) was added 1N NaOH aqueous solution (1ml) followed by dropwise addition of 3, 4-dichlorophenylacetyl chloride at room temperature. After 2.5h stirring, the reaction mixture was extracted with CH<sub>2</sub>Cl<sub>2</sub>. The extract combined was washed with brine, dried (MgSO<sub>4</sub>), and concentrated to give 0.952g of brown viscous oil, which was purified by column chromatography (silica gel : 30g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH : 30/1 to 10/1 as eluent) to afford 152mg (28.1%) of title compound as yellow brown viscous oil.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.39-7.27 (11H, m), 7.15-7.08 (1.5H, m), 7.01 (0.5H, dd, J = 2.2, 8.4Hz), 6.04 (0.5H, dd, J = 5.9, 9.9Hz), 5.10 (1H, s), 5.07 (0.5H, d, J = 12.1Hz), 5.00 (0.5H, t, J = 7.5Hz), 4.99 (0.5H, d, J = 12.1Hz), 4.28 (0.5H, d, J = 16.8Hz), 4.30-4.20 (1H, m), 3.97 (0.5H, d, J = 18.7Hz), 3.92 (0.5H, d, J = 16.5Hz), 3.89 (0.5H, d, J = 15.7Hz), 3.76

-13-

(1H, s), 3.71 (0.5H, d, J = 16.1Hz), 3.60 (0.5H, d, J = 15.7Hz), 3.14 - 2.42 (5H, m), 2.27 - 2.01 (2H, m), 1.80 - 1.60 (2H, m).

IR (neat) : 3450, 1750, 1650 cm<sup>-1</sup>.

This oil was converted to HCl salt using HCl gas saturated methanol to give 77mg of amorphous  
5 solid.

Anal. Calcd for C<sub>28</sub>H<sub>30</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>4</sub>·HCl·1.5H<sub>2</sub>O : C, 57.58 ; 5.66; N, 4.63 Cl, 17.58.

Found : C, 57.56 ; H, 5.37 ; N, 4.71 ; Cl, 17.67.

#### Example 2

(2S,3S)-1-[2-N-(t-Butoxycarbonyl) methylamino-2-phenylethyl]-3-hydroxypyrrolidine

10 This compound was prepared in 70% yield according to a procedure similar to that described in Preparation 3.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.34-7.28 (5H, m), 4.38-4.30 (1H, m), 3.81 (1H, dd, J = 3.7, 10.6Hz), 3.25 (1H, d, J = 17.6Hz), 3.19-3.13 (1H, m), 3.04 (1H, d, J = 17.6Hz), 2.87 (1H, dd, J = 10.6, 12.1Hz), 2.75-2.65 (2H, m), 2.34 (1H, dd, J = 3.7, 11.7Hz), 2.27-2.15 (4H, m),

15 1.85-1.75 (1H, m), 1.44 (9H, s).

IR(neat) : 3300, 1730cm<sup>-1</sup>.

#### Example 3

N-(t-Butoxycarbonyl)methyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

20 This compound was prepared in 37.1% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.41-7.10 (8H, m), 6.10 (0.7H, dd, J = 6.6, 9.6Hz), 5.03 (0.3H, dd, J = 7.0, 7.7Hz), 4.30-4.20 (1H, m), 3.93 (0.3H, d, J = 16.8Hz), 3.82 (0.6H, s), 3.82 (0.7H, d, J = 18.7Hz), 3.76 (0.3H, d, J = 15.4Hz), 3.71 (0.7H, d, J = 18.7Hz), 3.62 (0.7H, d, J = 15.8Hz), 3.55 (0.7H, d, J = 15.8Hz), 3.17-2.05 (9H, m), 1.80-1.65 (1H, m), 1.39 (2.7H, s), 1.30 (6.3H, s).

IR(neat) : 3400, 1740, 1650cm<sup>-1</sup>.

Anal. Calcd for C<sub>28</sub>H<sub>22</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>4</sub>·0.5H<sub>2</sub>O : C, 60.47 ; H, 6.44 ; N, 5.42.

Found : C, 60.24 ; H, 6.41 ; N, 5.24.

30

#### Example 4

N-Carboxymethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

A mixture of N-(t-butoxycarbonyl)methyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide (306mg, 0.6mmol), HCl gas saturated methanol solution (8ml),



-14-

and methanol (2ml) was refluxed with stirring for 2h. After evaporation of the solvent, the resulting oil was crystallized from  $\text{CH}_2\text{Cl}_2/\text{MeOH}$  to give 153mg (52.3%) of desired compound as HCl salt of white powder.

mp 161.9 - 163.5 °C

5 Anal. Calcd for  $\text{C}_{22}\text{H}_{24}\text{Cl}_2\text{N}_2\text{O}_4 \cdot \text{HCl}$  : C, 54.17 ; H, 5.17 ; N, 5.74.

Found : C, 54.40 ; H, 5.47 ; N, 5.62.

$^1\text{H}$ NMR (270MHz,  $\text{CDCl}_3$ )  $\delta$  11.30 (1H, br.s), 7.45-7.25 (8H, m), 6.45 (1H, m), 4.65-2.00 (15H, m).

IR (KBr) : 3300, 1740, 1665 $\text{cm}^{-1}$ .

10

#### Example 5

##### (2S,3S)-1-[2-N-(2-Hydroxyethylamino)-2-phenylethyl]-3-hydroxypyrrolidine

To a stirred suspension of lithium aluminum hydride (380mg, 10mmol) in THF (10ml) was added a solution of (2S, 3S)-1-[2-N-(t-butoxycarbonyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine (1.602g, 5mmol) in THF (15ml) dropwise at room temperature. The reaction mixture was then refluxed with stirring for 1h. After cooling down to room temperature,  $\text{Na}_2\text{SO}_4 \cdot 10\text{H}_2\text{O}$  (3.80g) and KF (0.38g) was added to the reaction mixture. After 1h stirring, the solid appeared was removed by Celite filtration. The filtrate was concentrated to give 1.32g (crude 100%) of desired compound as yellow oil.

15  $^1\text{H}$  NMR (270MHz,  $\text{CDCl}_3$ )  $\delta$  7.34-7.24 (5H, m), 4.33 (1H, br.s), 3.77 (1H, dd, J = 3.7, 11.0Hz), 3.77-3.57 (2H, m), 3.20-3.05 (1H, m), 2.88 (1H, dd, J = 11.0, 12.1Hz), 3.05-2.55 (6H, m), 2.35 (1H, dd, J = 3.7, 12.1Hz), 2.30-2.15 (2H, m), 1.95-1.75 (2H, m).

IR (neat) : 3350 $\text{cm}^{-1}$ .

#### Example 6

##### 2-(3,4-Dichlorophenyl)-N-(2-hydroxyethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-

25 phenylethyl]acetamide

This compound was prepared in 48% yield according to a procedure similar to that described in Example 1.

$^1\text{H}$  NMR (270MHz,  $\text{CDCl}_3$ )  $\delta$  7.38-7.02 (8H, m), 6.13 (0.75H, br.d, J = 9.2Hz), 5.05 (0.25H, m), 4.45-4.35 (1H, m), 4.10-3.10 (9H, m), 2.90 -2.15 (6H, m), 1.90-1.75 (1H, m).

30 IR (neat) : 3400, 1640  $\text{cm}^{-1}$ .

HCl salt : amorphous solid.

Anal. Calcd for  $\text{C}_{22}\text{H}_{22}\text{Cl}_2\text{N}_2\text{O}_3 \cdot \text{HCl} \cdot 2.5\text{H}_2\text{O}$  : C, 50.93 ; H, 6.22 ; N, 5.40. Found : C, 51.20 ; H, 6.02 ; N, 5.66.

#### Example 7

(2S,3S)-3-Hydroxy-1-[2-phenyl-2-N-(2,2,2-trifluoroethylamino)ethyl]pyrrolidine

- To a stirred solution of (S)-phenylglycyl-3-(S)-hydroxypyrrolidine (1.00g, 4.5mmol) in THF (20ml) was added trifluoroacetic anhydride (0.7ml, 5mmol) at room temperature. After 1h stirring, the solvent was evaporated and the residue was purified by column chromatography (silica gel:100g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH:20/1 as eluent) to give 0.90g of amide derivative, which was dissolved in THF (14ml) followed by addition of boran-methyl sulfide complex (1.33ml, 14mmol) at 0°C. Then the reaction mixture was refluxed for 13h. To this reaction mixture was added 1N HCl aqueous solution (10ml) at 0°C and the mixture was refluxed for 1h. After cooling down to room temperature, the reaction mixture was basified with 1N NaOH aqueous solution and extracted with CH<sub>2</sub>Cl<sub>2</sub>. After dry (Na<sub>2</sub>SO<sub>4</sub>), the solvent was evaporated to give 0.821g of colorless oil which was soon crystallized.
- <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.38-7.20 (5H, m), 4.38-4.30 (1H, m), 3.92 (1H, dd, J = 3.3, 11.0Hz), 3.13-2.95 (3H, m), 2.81 (1H, dd, J = 11.0, 12.1Hz), 2.85-2.60 (3H, m), 2.38-2.11 (3H, m), 1.95 (1H, br.s), 1.85-1.65 (1H, m).
- IR(neat) : 3350cm<sup>-1</sup>.

Example 82-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(2,2,2-trifluoroethyl)acetamide

- This compound was prepared in 38.9% yield according to a procedure similar to that described in Example 1.
- <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.40 (1H, d, J = 8.1Hz), 7.36-7.26 (5H, m), 7.16 (1H, br.s), 7.07 (1H, br.d, J = 8.1Hz), 5.78 and 5.06 (total 1H, each br.s), 4.27 (1.5H, br.s), 3.90-3.60 (3.5H, m), 3.80-2.80 (3H, m), 2.75-2.60 (2H, m), 2.45-2.05 (2H, m), 1.95-1.60 (2H, m).
- IR (neat) : 3450, 1660 cm<sup>-1</sup>.
- HCl salt : amorphous solid.
- Anal. Calcd for C<sub>22</sub>H<sub>22</sub>Cl<sub>2</sub>F<sub>3</sub>N<sub>2</sub>O<sub>2</sub>·HCl : C, 51.63 ; H, 4.73 ; N, 5.47.
- Found : C, 51.78 ; H, 5.19 ; N, 5.28.

Example 9(2S,3S)-1-[2-N-(2-furyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine

- A mixture of (2S,3S)-1-(2-amino-2-phenylethyl)-3-hydroxypyrrolidine (619mg, 3mmol), and 2-furaldehyde (0.37ml, 4.5mmol) in ethanol (8ml) was refluxed with stirring for 4.5h. Then the solvent was evaporated and the resulting Schiff base was dissolved in methanol (10ml) and to this solution was added NaBH<sub>4</sub> by portions at room temperature. After 0.5h stirring at room temperature, the solvent was evaporated. Then water (30ml) was added to the residue and

-16-

extracted with  $\text{CH}_2\text{Cl}_2$ . The extract was washed with brine, dried ( $\text{MgSO}_4$ ), and concentrated to give 857mg (99.8%) of brown oil.

$^1\text{H}$  NMR (270MHz,  $\text{CDCl}_3$ )  $\delta$  7.40-7.28 (6H, m), 6.30 (1H, dd,  $J = 1.8, 2.9\text{Hz}$ ), 6.08 (1H, d,  $J = 2.9\text{Hz}$ ), 4.33-4.25 (1H, m), 3.75 (1H, d,  $J = 14.7\text{Hz}$ ), 3.71 (1H, dd,  $J = 3.7, 11.4\text{Hz}$ ),  
5 3.49 (1H, d,  $J = 14.7\text{Hz}$ ), 2.84 (1H, dd,  $J = 11.4, 12.1\text{Hz}$ ), 2.83-2.75 (1H, m), 2.62-2.51 (2H, m), 2.35 (2H, br.s), 2.27 (1H, dd,  $J = 3.7, 12.1\text{Hz}$ ), 2.21-2.10 (2H, m), 1.80-1.65 (1H, m).  
IR (neat) :  $3300\text{cm}^{-1}$ .

#### Example 10

##### 2-(3,4-Dichlorophenyl)-N-furfuryl-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-

##### phenylethyl]acetamide

This compound was prepared in 96.5% yield according to a procedure similar to that described in Example 1.

$^1\text{H}$  NMR (270MHz,  $\text{CDCl}_3$ )  $\delta$  7.37 (1H, d,  $J = 8.1\text{Hz}$ ), 7.37 (1H, d,  $J = 1.8\text{Hz}$ ), 7.31-7.22 (5H, m), 7.10 (1H, dd,  $J = 1.8, 8.1\text{Hz}$ ), 7.03 (1H, dd,  $J = 1.8, 8.1\text{Hz}$ ), 6.19 (1H, dd,  $J = 1.9, 2.9\text{Hz}$ ), 5.94 (1H, dd,  $J = 5.1, 9.5\text{Hz}$ ), 5.71 (1H, d,  $J = 2.9\text{Hz}$ ), 4.40-4.33 (1H, m), 4.30 (1H, d,  $J = 17.9\text{Hz}$ ), 4.22 (1H, d,  $J = 17.9\text{Hz}$ ), 3.72 (1H, d,  $J = 13.2\text{Hz}$ ), 3.66 (1H, d,  $J = 13.2\text{Hz}$ ), 3.59 (1H, dd,  $J = 9.5, 12.8\text{Hz}$ ), 3.49 (1H, s), 3.45-3.35 (1H, m), 3.20-3.05 (2H, m), 2.83 (1H, dd,  $J = 5.1, 11.0\text{Hz}$ ), 2.65-2.55 (1H, m), 2.55-2.10 (1H, m), 1.95-1.80 (1H, m).  
IR (neat) : 3400,  $1650\text{cm}^{-1}$ .

20 HCl salt : mp  $181.0 - 183.5^\circ\text{C}$

Anal. Calcd for  $\text{C}_{25}\text{H}_{26}\text{Cl}_2\text{N}_2\text{O}_3$  : C, 58.89 ; H, 5.34 ; N, 5.49.

Found : C, 58.58 ; H, 5.61 ; N, 5.63.

#### Example 11

##### (2S,3S)-1-[2-Phenylethyl-2-N-(2-thienyl)methylamino]-3-hydroxypyrrolidine

25 This compound was prepared in 100% yield according to a procedure similar to that described in Example 9.

$^1\text{H}$  NMR (270MHz,  $\text{CDCl}_3$ )  $\delta$  7.42-7.24 (5H, m), 7.20 (1H, dd,  $J = 1.1, 5.1\text{Hz}$ ), 6.94 (1H, dd,  $J = 3.3, 5.1\text{Hz}$ ), 6.83 (1H, br.d,  $J = 2.9\text{Hz}$ ), 4.30-4.20 (1H, m), 3.90 (1H, d,  $J = 14.3\text{Hz}$ ), 3.79 (1H, dd,  $J = 3.7, 11.0\text{Hz}$ ), 3.73 (1H, d,  $J = 14.3\text{Hz}$ ), 2.84 (1H, dd,  $J = 11.0, 12.1\text{Hz}$ ),  
30 2.88-2.77 (1H, m), 2.62-2.52 (2H, m), 2.29 (1H, dd,  $J = 3.7, 12.1\text{Hz}$ ), 2.32-2.10 (4H, m), 1.77-1.64 (1H, m).

IR (neat) :  $3300\text{cm}^{-1}$ .

#### Example 12

-17-

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(2-thienyl)methylacetamide

This compound was prepared in 62.6% yield according to a procedure similar to that described in Example 1.

- 5 <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.39 (1H, d, J = 2.2Hz), 7.33 (1H, d, J = 8.8Hz), 7.30-7.26 (3H, m), 7.15-7.09 (3H, m), 6.94 (1H, dd, J = 2.2, 8.4Hz), 6.83 (1H, dd, J = 3.7, 5.1Hz), 6.61 (1H, br.d, J = 2.9Hz), 5.99 (1H, dd, J = 5.1, 9.5Hz), 4.59 (1H, d, J = 17.9Hz), 4.47 (1H, d, J = 17.6Hz), 4.43-4.35 (1H, m), 3.67-3.35 (5H, m), 3.21-3.08 (2H, m), 2.87 (1H, dd, J = 5.1, 11.0Hz), 2.68-2.56 (1H, m), 2.26-2.14 (1H, m), 1.96-1.85 (1H, m).
- 10 IR (neat) : 3400, 1650cm<sup>-1</sup>.  
HCl salt : mp 180.7 - 184.0°C  
Anal. Calcd for C<sub>25</sub>H<sub>26</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>2</sub>S·HCl·0.4H<sub>2</sub>O : C, 56.32 ; H, 5.26 ; N, 5.25.  
Found : C, 56.67 ; H, 5.38 ; N, 4.77.

Example 13

- 15 (2S,3S)-1-[2-Phenylethyl-2-N-(3-pyridyl)methylaminol]-3-hydroxypyrrolidine

This compound was prepared in 63.9% yield according to a procedure similar to that described in Example 9.

- <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 8.49(1H, s), 8.48 (1H, d, J = 4.8Hz), 7.63 (1H, dd, J = 1.5, 7.7Hz), 7.42-7.27 (5H, m), 7.24 (1H, dd, J = 4.8, 7.7Hz), 4.35-4.25 (1H, m), 3.74 (1H, dd, J = 3.7, 11.0Hz), 3.73 (1H, d, J = 13.8Hz), 3.55 (1H, d, J = 13.6Hz), 2.85 (1H, dd, J = 11.0, 12.1Hz), 2.88-2.82 (1H, m), 2.60 (2H, d, J = 4.0Hz), 2.42 (2H, br.s), 2.31 (1H, dd, J = 3.7, 12.1Hz), 2.27-2.09 (2H, m), 1.80-1.65 (1H, m).
- 20 IR (neat) : 3300cm<sup>-1</sup>.

Example 14

- 25 2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(3-pyridyl)methylacetamide

This compound was prepared in 56.7% yield according to a procedure similar to that described in Example 1.

- <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 8.51 (0.7H, s), 8.48 (0.7H, d, J = 4.4Hz), 8.40 (0.3H, d, J = 2.9Hz), 8.26 (0.3H, s), 7.55-6.99 (10H, m), 6.26 (0.7H, dd, J = 4.0, 11.0Hz), 5.17 (0.3H, t), 4.70 (0.3H, d), 4.41-4.20 (2.7H, m), 3.90 (0.6H, s), 3.58 (0.7H, d), 3.50(0.7H, d), 3.25-2.35 (5H, m), 2.28-1.95 (2H, m), 1.85-1.60 (2H, m).
- 30 IR (neat) : 3300, 1650 cm<sup>-1</sup>.  
Anal. Calcd for C<sub>28</sub>H<sub>27</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub>·2HCl·1.5H<sub>2</sub>O : C, 53.44 ; H, 5.52 ; N, 7.19.

-18-

Found : C, 53.17 ; H, 5.21 ; N, 6.91.

#### Example 15

##### (2S,3S)-1-[2-Phenylethyl-2-N-(2-pyridyl)methylamino]-3-hydroxypyrrolidine

This compound was prepared in 37.9% yield according to a procedure similar to that described in Example 9.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 8.56 (1H, br.d, J = 4.8Hz), 7.61 (1H, dt, J = 1.8, 7.7Hz), 7.42-7.13 (7H, m), 4.33-4.25 (1H, m), 3.83 (1H, d, J = 14.3Hz), 3.73 (1H, dd, J = 3.7, 10.6Hz), 3.67 (1H, d, J = 14.3Hz), 3.00-2.80 (3H, m), 2.92 (1H, dd, J = 10.6, 12.1Hz), 2.70 (1H, br.d, J = 9.5Hz), 2.59 (1H, dd, J = 4.4, 9.9Hz), 2.36 (1H, dd, J = 3.7, 12.1Hz), 2.25-2.09 (2H, m), 1.85-1.70 (1H, m).

IR (neat) : 3300cm<sup>-1</sup>.

#### Example 16

##### 2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(2-pyridyl)methylacetamide

This compound was prepared in 68.5% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 8.79 (1H, br.s), 7.66-7.60 (1H, m), 7.33 (1H, d, J = 8.4Hz), 7.33-7.23 (6H, m), 7.18 (1H, d, J = 1.8Hz), 7.05 (1H, br.d, J = 7.7Hz), 6.97 (1H, dd, J = 1.8Hz, 8.1Hz), 6.50 (1H, dd, J = 3.7, 12.1Hz), 4.70-4.55 (2H, m), 4.37 (1H, d, J = 18.3Hz), 4.20 (1H, dd, J = 12.5, 12.8Hz), 3.95-3.25 (8H, m), 2.50-2.30 (1H, m), 2.25-2.10 (1H, m).

IR (neat) : 3400, 1650cm<sup>-1</sup>.

HCl Salt : mp 126.5 - 132.0 °C

Anal. Calcd for C<sub>26</sub>H<sub>27</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub>·2HCl·2H<sub>2</sub>O : C, 52.63 ; H, 5.61 ; N, 7.08

Found : C, 52.31 ; H, 5.39 ; N, 6.75.

25

#### Example 17

##### (2S,3S)-1-[2-Phenylethyl-2-N-(4-pyridyl)methylamino]-3-hydroxypyrrolidine

This compound was prepared in 81.8% yield according to a procedure similar to that described in Example 9.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 8.52 (2H, d, J = 5.5Hz), 7.40-7.25 (5H, m), 7.22 (2H, d, J = 5.5Hz), 4.37-4.27 (1H, m), 3.72 (1H, d, J = 14.3Hz), 3.75-3.70 (1H, m), 3.56 (1H, d, J = 14.7Hz), 2.86 (1H, t, J = 11.4Hz), 2.90-2.82 (1H, m), 2.62 (2H, app d), 2.39 (2H, br.s), 2.35-2.10 (3H, m), 1.85-1.68 (1H, m).

IR (neat) : 3300cm<sup>-1</sup>.

#### Example 18

2-(3,4-Dichlorophenyl)-N-[2-(3-(s)-hydroxypyrrolidin-1-yl)-1-(s)-phenylethyl]-N-(4-pyridyl)methylacetamide

This compound was prepared in 10.9% yield according to a procedure similar to that described in Example 1.

- 5 <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 8.48 (1.4H, d, J = 5.9Hz), 8.40 (0.6H, d, J = 5.9Hz), 7.44-6.97 (10H, m), 6.26 (0.7H, dd, J = 4.4, 11.0Hz), 5.17 (0.3H, app t), 4.62 (0.3H, d, J = 16.1Hz), 4.40-4.30 (1H, m), 4.29 (1.4H, s), 4.26 (0.3H, d, J = 16.9Hz), 3.92 (0.6H, s), 3.49 (0.7H, d, J = 15.4Hz), 3.41 (0.7H, d, J = 15.4Hz), 3.20-2.45 (5H, m), 2.35-2.00 (2H, m), 1.92 (1H, br.s), 1.85-1.65 (1H, m).

- 10 IR (neat) : 3400, 1650cm<sup>-1</sup>.

HCl salt : mp 161.1 - 164.2

Anal. Calcd for C<sub>26</sub>H<sub>27</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub>·2HCl·3H<sub>2</sub>O : C, 51.08 ; H, 5.77 ; N, 6.87.

Found : C, 50.75 ; H, 5.26 ; N, 6.83.

**Example 19**

- 15 (2S,3S)-1-[2-N-(4-Methoxycarbonylphenyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine

This compound was prepared in 81.3% yield according to a procedure similar to that described in Example 9.

- <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.99 (2H, d, J = 8.1Hz), 7.40-7.29 (7H, m), 4.35-4.25 (1H, m), 3.91 (3H, s), 3.79 (1H, d, J = 13.9Hz), 3.70 (1H, dd, J = 3.3, 10.6Hz), 3.57 (1H, d, J = 13.9Hz), 2.85 (1H, dd, J = 10.6, 12.1Hz), 2.85-2.81 (1H, m), 2.62-2.52 (2H, m), 2.30 (1H, dd, J = 3.3, 12.1Hz), 2.27-2.00 (4H, m), 1.80-1.65 (1H, m).

IR (neat) : 3280, 3200, 1720cm<sup>-1</sup>.

**Example 20**

- 25 2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(4-methoxycarbonylphenyl)methylacetamide

This compound was prepared in 78.5% yield according to a procedure similar to that described in Example 1.

- <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.83 (2H, d, J = 8.1Hz), 7.39-6.90 (10H, m), 6.30-6.15 (1H, m), 4.60-4.10 (3H, m), 3.90 (3H, s), 3.68 (1H, t, J = 11.7Hz), 3.63-3.45 (2H, m), 3.30-3.17 (2H, m), 3.10-2.95 (2H, m), 2.74 (1H, br.s), 2.35-2.20 (1H, m), 2.05-1.90 (1H, m).

IR (neat) : 3400, 1720, 1650cm<sup>-1</sup>.

HCl salt : amorphous solid

Anal. Calcd for C<sub>29</sub>H<sub>31</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>4</sub>·HCl·0.7H<sub>2</sub>O : C, 58.88 ; H, 5.69 ; N, 4.74.

Found : C, 58.56 ; H, 5.24 ; N, 4.65.

-20-

**Example 21****(2S,3S)-1-(2-N-Cyanomethylamino-2-phenylethyl)-3-hydroxypyrrolidine**

To a stirred solution of sodium bisulfite (1.249g, 12mmol) in water (1ml) was added 37% aqueous solution of formaldehyde (0.9ml, 12mmol) at 0 and the mixture was stirred at room temperature for 0.5h. Then to this mixture was added (2S,3S)-1-(2-phenylethyl)-3-hydroxypyrrolidine (2.48g, 12mmol) and the resulting mixture was stirred at 50 for 0.5h. After cooling down to room temperature, aqueous solution of KCN(781.4mg, 12mmol) was added to the reaction mixture and stirring was continued for 2h. The reaction mixture was diluted with water (30ml), extracted with CH<sub>2</sub>Cl<sub>2</sub> (20ml x 3). The extract was washed with saturated NaHCO<sub>3</sub> aqueous solution and dried (Na<sub>2</sub>SO<sub>4</sub>). The solvent was evaporated to give 2.20g of yellow oil, which was purified by column chromatography (silica gel ; 70g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH : 50/1 as eluent) to afford 1.725g (58.6%) of colorless viscous oil which was gradually crystallized.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.39-7.27 (5H, m), 4.40-4.35 (1H, m), 3.96 (1H, dd, J = 3.3, 11.4Hz), 3.67 (1H, d, J = 17.6Hz), 3.24 (1H, d, J = 17.6Hz), 3.17-3.09 (1H, m), 2.87 (1H, app t, J = 11.7Hz), 2.75 (1H, dd, J = 4.8, 9.9Hz), 2.67 (1H, br.d, J = 8.4Hz), 2.36 (1H, dd, J = 3.3, 12.1Hz), 2.31-2.10 (3H, m), 1.90 (1H, br.s), 1.88 -1.75(1H, m).

IR (neat) : 3300, 2230cm<sup>-1</sup>.

**Example 22****N-Cyanomethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide**

This compound was prepared in 46.6% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.38-7.00 (8H, m), 6.09 (0.7H, br.s), 5.31 (0.3H, br.s), 5.09 (1H, br.s), 4.45 (1H, Br.d, J = 16.9Hz), 4.40-4.25 (1H, m), 4.15-3.70 (4H, m, including 1H, br.d, J = 16.1Hz at 3.95ppm), 3.65-2.55 (4H, m), 2.50-1.70 (4H, m).

<sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 171.2, 137.4, 134.0, 132.7, 131.4, 131.1, 130.6,

129.3, 128.6, 127.3, 126.3, 116.7, 71.2, 63.2, 59.9, 57.9, 51.9, 39.4, 35.0, 31.0.

IR (neat) : 3450, 2250, 1660cm<sup>-1</sup>.

MS m/e (%) : 432 (<4), 189 (22), 161 (93), 159 (100), 145 (39), 143 (35), 132 (32), 125 (31), 123 (37), 117 (30), 100 (99).

**Preparation 4****3-(S)-Methoxymethoxypyrrolidine**

To a stirred solution of N-benzyl-3-(S)-hydroxypyrrolidine (4.785g, 27mmol) in THF (50ml) was added sodium hydride (60% oil suspension, 1.12g, 28mmol) by portions under nitrogen at room

-21-

temperature. The suspension mixture was then refluxed for 1h. To this reaction mixture was added chloromethyl methyl ether (2.3ml, 30mmol) and refluxing was continued for 13h. After cooling down to room temperature, water (10ml) was added to the reaction mixture to give a solution mixture, which was basified with 1N NaOH aqueous solution, extracted with ethyl acetate (30ml x 3), and dried (Na<sub>2</sub>SO<sub>4</sub>). Evaporation of the solvent gave 6.33g of brown oil, which was purified by column chromatography (silica gel : 150g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH : 20/1 as eluent) to afford 5.09g (85%) of clear brown oil. A suspension mixture of this oil (5.09g, 23mmol) and Pearlman's catalyst (2.00g) in MeOH (100ml) was stirred under hydrogen atmosphere at room temperature for 15h. After removal of the catalyst by Celite filtration, the filtrate was concentrated to give an oil and water mixture, which was dried in vacuo to afford 2.76g (91.4%) of orange color oil.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 4.65 (1H, d, J = 7.0Hz), 4.63 (1H, d, J = 6.6Hz), 4.30-4.24 (1H, m), 3.37 (3H, s), 3.21 (1H, br.s), 3.17-2.86 (4H, m), 1.98-1.80 (2H, m).  
IR(neat) : 3300cm<sup>-1</sup>.

15

#### Preparation 5

##### (S)-Phenylglycyl-3-(S)-methoxymethoxypyrrolidine

This compound was prepared in 61.8% yield using 3-(S)-methoxymethoxypyrrolidine and N-benzyloxycarbonyl-(S)-phenylglycine according to a procedure similar to that described in Preparation 1.

20 <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.36-7.29 (5H, m), 4.66 and 4.63, and 4.46 and 4.35 (total 2H, each d, J = 7.0Hz), 4.56 and 4.50 (total 1H, each br.s), 4.27-4.20 (1H, m), 3.81-3.14 (4H, m), 3.36 and 3.10 (total 3H, each s), 2.10-1.80 (2H, m), 2.00 (2H, br.s).  
IR (neat) : 3350, 3300, 1650 cm<sup>-1</sup>.

#### Preparation 6

##### (2S,3S)-1-(2-Amino-2-phenylethyl)-3-(methoxymethoxy)pyrrolidine

This compound was prepared in 97.7% yield according to a procedure similar to that described in Preparation 2.

30 <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.39-7.24 (5H, m), 4.66 (1H, d, J = 6.6Hz), 4.63 (1H, d, J = 7.0Hz), 4.30-4.23 (1H, m), 4.08 (1H, dd, J = 3.3, 10.3Hz), 3.37 (3H, s), 3.00-2.80 (2H, m), 2.74 (1H, dd, J = 10.3, 11.7Hz), 2.60-2.45 (2H, m), 2.41 (1H, dd, J = 3.3, 11.7Hz), 2.21-2.04 (1H, m), 1.95-1.75 (3H, m).  
IR (neat) : 3400 cm<sup>-1</sup>.

#### Example 23

##### (2S,3S)-1-[2-N-(2,2-Difluoroethylamino)-2-phenylethyl]-3-hydroxypyrrolidine



-22-

T a stirred solution of (2S,3S)-1-(2-amino-2-phenylethyl)-3-(methoxymethoxy)pyrrolidine (0.77g, 3.08mmol) and 2,2-difluoroacetic acid (0.21ml, 3.3mmol) was added 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.63g, 3.3mmol) at room temperature. After 4h stirring, the reaction mixture was diluted with CH<sub>2</sub>Cl<sub>2</sub> (20ml), washed with saturated NaHCO<sub>3</sub> aqueous solution and brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to give 1.05g of brown viscous oil. This was purified by column chromatography (silica gel: 100g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH : 40/1 as eluent) to give 0.69g (68.3%) of brown viscous oil.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.37-7.25 (5H, m), 5.91 (1H, t, J<sub>H,F</sub> = 54.2Hz), 4.95-4.85 (1H, m), 4.63 (1H, d, J = 6.6Hz), 4.59 (1H, d, J = 6.6Hz), 4.26-4.19 (1H, m), 3.35 (3H, s), 2.96-2.69 (4H, m), 2.56-2.43 (2H, m), 2.18-2.03 (1H, m), 1.87-1.75 (1H, m), 1.72 (1H, br.s). IR (neat) : 3300, 1700cm<sup>-1</sup>.

To a stirred solution of this amide derivative (0.69g, 2.1mmol) in THF (6ml) was added BH<sub>3</sub>·Me<sub>2</sub>S (0.6ml, 6.3mmol) at room temperature and then refluxed for 5h. After cooling down to room temperature, 1N HCl (10ml) was added dropwise carefully and the resulting mixture was refluxed for 2h. After cooling down to room temperature, the reaction mixture was basified with 1N NaOH aqueous solution to pH 12 and extracted with CH<sub>2</sub>Cl<sub>2</sub> (20ml x 3). The extract was dried (Na<sub>2</sub>SO<sub>4</sub>) and concentrated to give 566mg (67.9% for two steps) of titled compound as yellow viscous oil.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.37-7.23 (5H, m), 5.76 (1H, ddt, J = 3.3, 5.1, 56.1Hz), 4.40-4.25 (1H, m), 3.80 (1H, dd, J = 3.7, 10.6Hz), 3.77-3.70 (1H, m), 2.99 (1H, dt, J = 5.5, 8.8Hz), 2.95-2.45 (4H, m), 2.82 (1H, dd, J = 11.0, 12.1Hz), 2.36-1.95 (3H, m), 1.90-1.70 (2H, m). IR (neat) : 3400, 3330cm<sup>-1</sup>.

#### Example 24

##### 25 2-(3,4-Dichlorophenyl)-N-(2,2-difluoroethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This compound was prepared in 62.7% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.43-7.26 (6H, m), 7.15-7.09 (2H, m), 6.13(0.5H, dd, J = 5.5, 11.0Hz), 6.03 (0.5H, ddt, J = 2.9, 5.1, 55.0Hz), 5.27 (0.5H, ddt, J = 4.4, 4.8, 56.4Hz), 5.05(0.5H, dd, J = 6.6, 7.7Hz), 4.35-4.25 (1H, m), 3.86 (1H, s), 3.84 (0.5H, d, J = 15.8Hz), 3.75(0.5H, d, J = 15.8Hz), 3.70-2.60 (7H, m), 1.80-1.65 (1H, m).

IR (neat) : 3450, 1650cm<sup>-1</sup>.

HCl salt : amorphous solid

-23-

Anal. Calcd for  $C_{22}H_{24}Cl_2F_2N_2O_2 \cdot HCl \cdot H_2O$  : C, 51.63 ; H, 5.32 ; N, 5.47.

Found : C, 51.94 ; H, 5.40 ; N, 5.51.

#### Preparation 7

##### (2S,3S)-1-[2-N-(2-Fluoroethylamino)-2-phenylethyl]-3-(methoxymethoxy)pyrrolidine

- 5 A mixture of (2S,3S)-1-(2-amino-2-phenylethyl)-3-(methoxymethoxy)pyrrolidine (1.01g, 4mmol), 2-bromofluoroethane (0.90ml, 12mmol), and  $K_2CO_3$  (0.69g, 5mmol) in DMF (5ml) was stirred at 70°C for 13h. The reaction mixture was diluted with water (10ml), basified with 1N NaOH aqueous solution to pH 12, and extracted with  $CH_2Cl_2$  (30ml x 3). The extract was dried ( $Na_2SO_4$ ) and concentrated in vacuo to give 1.00g of dark brown viscous oil, which was purified by column
- 10 chromatography (silica gel: 100g,  $CH_2Cl_2$ /MeOH : 30/1 as eluent) to afford 0.51g (43%) of brown viscous oil.

$^1H$  NMR (270MHz,  $CDCl_3$ )  $\delta$  7.38-7.25 (5H, m), 4.66 (1H, d,  $J = 7.0$ Hz), 4.63 (1H, d,  $J = 7.0$ Hz), 4.60-4.22 (3H, m), 3.77 (1H, dd,  $J = 3.3, 10.3$ Hz), 3.38 (3H, s), 2.98 (1H, dd,  $J = 6.2, 10.3$ Hz), 2.94-2.49 (6H, m), 2.34 (1H, dd,  $J = 3.7, 12.1$ Hz), 2.20-2.06 (1H, m), 1.90-1.78

15 (1H, m), 1.56 (1H, br.s).  
IR (neat) : 3320  $cm^{-1}$ .

#### Example 25

##### 2-(3,4-Dichlorophenyl)-N-(2-fluoroethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

- 20 To a stirred solution of (2S,3S)-1-[2-N-(2-fluoroethylamino)-2-phenylethyl]-3-(methoxymethoxy)pyrrolidine (0.7g, 2.6mmol) and triethylamine (0.56ml, 4mmol) in  $CH_2Cl_2$  (10ml) was added 3,4-dichlorophenylacetyl chloride [this was prepared from 3,4-dichlorophenylacetic acid (0.82g, 4mmol) and thionyl chloride (0.36ml, 5mmol)] at room temperature. After 20min, the reaction mixture was diluted with  $CH_2Cl_2$  (20ml), washed with
- 25 saturated  $NaHCO_3$  aqueous solution, and dried ( $Na_2SO_4$ ). Evaporation of the solvent gave 1.96g of brown oil, which was purified by column chromatography (silica gel 100g,  $CH_2Cl_2$ /MeOH : 20/1 as eluent) to give 0.81g (64.3%) of yellow viscous oil. To this oil was added HCl gas saturated MeOH solution (10ml) and stirred at room temperature for 2h. The solvent was
- 30 evaporated and the residue was basified with 1N NaOH aqueous solution to pH 11 and extracted with  $CH_2Cl_2$  (20ml x 2). After dry ( $Na_2SO_4$ ), the solvent was evaporated to afford 0.78g of brown viscous oil, which was purified by column chromatography (silica gel : 100g,  $CH_2Cl_2$ /MeOH : 40/1 as eluent) to give 0.598g (81.5%) of clear yellow viscous oil.
- $^1H$  NMR (270MHz,  $CDCl_3$ )  $\delta$  7.41-7.26 (6H, m), 7.20-7.09 (2H, m), 6.10 (0.6H, dd,  $J = 5.9, 10.6$ Hz), 5.03 (0.4H, t,  $J = 7.3$ Hz), 4.70-3.90 (3H, m), 3.83 (0.6H, d,  $J = 15.4$ Hz), 3.83

-24-

(0.8H, s), 3.75 (0.6H, d,  $J = 15.8\text{Hz}$ ), 3.72-3.25 (2H, m), 3.22-2.98 (2H, m), 2.94-2.80 (1H, m), 2.75 -2.08 (5H, m), 1.80-1.65 (1H, m).

IR (neat) : 3400, 1640 $\text{cm}^{-1}$ .

HCl salt : amorphous solid.

5 Anal. Calcd for  $\text{C}_{22}\text{H}_{25}\text{Cl}_2\text{FN}_2\text{O}_2 \cdot \text{HCl} \cdot 2\text{H}_2\text{O}$  : C, 51.63; H, 5.91; N, 5.47.

Found : C, 51.95; H, 5.64; N, 5.42.

#### Example 26

##### (2S,3S)-1-(2-N-Benzylamino-2-phenylethyl)-3-hydroxypyrrolidine

This compound was prepared in 100% yield according to a procedure similar to that described in Example 9.

$^1\text{H}$  NMR (270MHz,  $\text{CDCl}_3$ )  $\delta$  7.43-7.22 (10H, m), 4.30- 4.25 (1H, m), 3.77 (1H, d,  $J = 13.6\text{Hz}$ ), 3.73 (1H, dd,  $J = 3.7, 11.0\text{Hz}$ ), 3.49 (1H, d,  $J = 13.6\text{Hz}$ ), 2.87 (1H, dd,  $J = 11.0, 12.1\text{Hz}$ ), 2.85-2.75 (1H, m), 2.64-2.51 (2H, m), 2.38 (2H, br.s), 2.30 (1H, dd,  $J = 3.7, 12.1\text{Hz}$ ), 2.23-2.07 (2H, m), 1.77-1.65 (1H, m).

15 IR (neat) : 3300 $\text{cm}^{-1}$ .

#### Example 27

##### N-Benzyl-2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This compound was prepared in 62.6% yield according to a procedure similar to that described in Example 1.

$^1\text{H}$  NMR (270MHz,  $\text{CDCl}_3$ )  $\delta$  7.40-6.78 (13H, m), 6.27 (1H, br.d,  $J = 8.1\text{Hz}$ ), 4.93 (1H, br.d,  $J = 17.6\text{Hz}$ ), 4.65-4.50 (2H, m), 4.20-3.95 (2H, m), 3.85-3.60 (3H, m), 3.55-3.15 (4H, m), 2.50-2.35 (1H, m), 2.30-2.10 (1H, m).

IR (KBr) : 3330, 1640 $\text{cm}^{-1}$ .

25 HCl salt : amorphous solid

Anal. Calcd for  $\text{C}_{27}\text{H}_{28}\text{Cl}_2\text{N}_2\text{O}_2 \cdot \text{HCl} \cdot \text{H}_2\text{O}$  : C, 60.29 ; H, 5.81 ; N, 5.21.

Found : C, 60.49 ; H, 5.38 ; N, 5.24.

#### Example 28

##### N-Carbamoylmethyl-2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

A mixture of N-cyanomethyl-2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxy-pyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide(2.54g, 5.9mmol) and HCl gas saturated methanol(20ml) was stirred at room temperature for 0.5h. The reaction mixture was concentrated, basified with aqueous  $\text{NH}_3$  solution, and extracted with  $\text{CH}_2\text{Cl}_2$  (30ml x 3). After dry( $\text{Na}_2\text{SO}_4$ ), the solvent was evaporated to give

-25-

brown viscous oil, which was purified by column chromatography (silica gel:100g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH: 30/1 to 15/1 as eluent) to afford 1.25g(47.3%) of pale yellow viscous oil.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 10.14 (1H, br.s), 7.40-7.09 (8H, m), 6.31 (1H, dd, J = 3.3, 12.8Hz), 5.70 (1H, br.s), 4.42 (1H, m), 3.80-3.55 (4H, m), 3.50-3.40 (1H, m), 3.35 (1H, app t, J = 12.8Hz), 2.76 (1H, J = 10.3Hz), 2.75-2.64 (2H, m, including 1H, dd, J = 3.3, 12.8Hz at 2.67ppm), 2.30-2.15 (2H, m), 1.90-1.70 (2H, m).

<sup>13</sup>C NMR (CDCl<sub>3</sub>) δ 173.2, 172.1, 137.4, 134.5, 132.5, 131.1, 130.4, 128.94, 128.86, 128.6, 128.3, 127.1, 70.4, 63.1, 53.9, 53.4, 51.4, 47.7, 39.8, 34.7.

IR (KBr) : 3400, 1690, 1650cm<sup>-1</sup>.

10 MS m/e (%) : 450 (3), 431 (2), 429 (3), 408 (2), 406 (3), 363 (4), 243 (3), 189 (75), 163 (99), 161 (94), 159 (99), 132 (36), 125 (55), 118 (100), 104 (95), 101 (98), 91 (91), 89 (70), 82 (61).

HCl salt : amorphous solid

Anal. Calcd for C<sub>22</sub>H<sub>25</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>3</sub>·HCl·H<sub>2</sub>O : C, 52.34 ; H, 5.59 ; N, 8.32.

Found : C, 52.41 ; H, 5.38 ; N, 7.96.

15

### Example 29

#### (2S,3S)-3-Hydroxy-1-[2-N-(2-methoxyethylamino)-2-phenylethyl]pyrrolidine

To a stirred solution of (2S,3S)-1-(2-amino-2-phenylethyl)-3-hydroxypyrrolidine (0.619g, 3mmol) and 2-methoxyacetic acid (0.23ml, 3mmol) in CH<sub>2</sub>Cl<sub>2</sub>(10ml) was added 1-ethyl-3-(3-dimethylaminopropyl)carbodiimide hydrochloride (0.863g, 4.5mmol) at room temperature. After 20 0.5h stirring, the reaction mixture was diluted with water(50ml) and extracted with CH<sub>2</sub>Cl<sub>2</sub> (30ml x 3). The extract was washed with saturated NaHCO<sub>3</sub> aqueous solution and brine, dried (Na<sub>2</sub>SO<sub>4</sub>, and concentrated to give 0.711g of clear yellow viscous oil. This was purified by column chromatography (silica gel:40g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH : 30/1 as eluent) to give 0.44g (52.7%) of pale 25 yellow viscous oil.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.35-7.22 (6H, m), 5.07-4.99 (1H, m), 4.30-4.20 (1H, m), 3.91 (2H, s), 3.42 (3H, s), 3.02 (1H, s), 2.94-2.85 (2H, m), 2.67 (1H, dd, J=5.1, 12.5Hz), 2.60 (2H, d, J=4.4Hz), 2.36-2.27 (1H, m), 2.17-2.07 (1H, m), 1.71-1.66 (1H, m).

IR (neat) : 3350, 3270, 1660cm<sup>-1</sup>.

30 To a stirred suspension of lithium aluminum hydride (0.24g, 6mmol) in THF (10ml) was added a solution of this amide derivative (0.44g, 1.58mmol) in THF(20ml) dropwise at room temperature and then the mixture was refluxed for 3.5h. After cooling down to room temperature, Na<sub>2</sub>SO<sub>4</sub> 10H<sub>2</sub>O(2.00g) and KF(0.2g) was added to the reaction mixture. After 20min stirring, the solid appeared was removed by Celite filtration. The filtrate was concentrated and the residue was

-26-

purified by column chromatography (silica gel:12g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH : 30/1 as eluent) to give 0.118g (28.2%) of pale yellow viscous oil.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.39-7.23 (5H, m), 4.35-4.26 (1H, m), 3.78 (1H, dd, J = 3.7, 10.6Hz), 3.53-3.40 (2H, m), 3.35 (3H, s), 3.10-3.00 (1H, m), 2.95-2.85 (3H, m), 2.87-2.60 (4H, m), 2.38 (1H, dd, J = 3.9, 12.1Hz), 2.33-2.15 (2H, m), 1.85-1.73 (1H, m).  
IR (neat) : 3400, 3330cm<sup>-1</sup>.

#### Example 30

##### 2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(2-methoxyethyl)acetamide

10 This compound was prepared in 77.2% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.40-7.25 (6H, m), 7.18-7.05 (2H, m), 5.93 (0.8H, dd, J = 5.9, 9.9Hz), 5.00 (0.2H, t), 4.35-4.25 (1H, m), 3.82 (1H, d, J = 15.4Hz), 3.75 (1H, d, J = 15.4Hz), 3.55-2.55 (13H, m, including 2.4H, s, at 3.19ppm), 2.50-2.35 (1H, m), 2.25 -2.10 (1H, m),  
15 1.85-1.70(1H, m).

IR (neat) : 3400, 1640cm<sup>-1</sup>.

HCl salt : amorphous solid.

Anal. Calcd for C<sub>22</sub>H<sub>22</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>3</sub>, HCl 2.2H<sub>2</sub>O : C, 52.37; H, 6.38; N, 5.31.

Found : C, 52.29; H, 6.40; N, 5.32.

20

#### Example 31

##### (2S,3S)-3-Hydroxy-1-[2-N-(2-methylthioethylamino)-2-phenylethyl]pyrrolidine

This compound was prepared in 50.5% yield according to a procedure similar to that described in Example 29.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.38-7.24 (5H, m), 4.35-4.26 (1H, m), 3.75 (1H, dd, J = 3.7, 10.6Hz), 3.10-3.00 (1H, m), 2.85 (1H, dd, J=10.6, 12.1Hz), 2.75-2.55 (5H, m), 2.41-2.15 (6H, m, including 1H, dd, J=4.0, 12.1Hz at 2.38ppm), 2.05 (3H, s), 1.85-1.70 (1H, m).

25 IR (neat) : 3300cm<sup>-1</sup>.

#### Example 32

##### 2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-

##### N-(2-methylthioethyl)acetamide

30 This compound was prepared in 64.8% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.45-7.25 (6H, m), 7.18-7.10 (2H, m), 6.06 (0.6H, dd, J = 5.9, 10.3Hz), 5.02 (0.4H, dd, J=6.2, 8.4Hz), 4.38-4.25 (1H, m), 3.83 (0.8H, s), 3.76 (0.6H, d,

-27-

J=15.4Hz), 3.68 (0.6H, d, J = 15.4Hz), 3.50-1.65 (16H, m, including 1.2H, s, at 1.99ppm and 1.8H, s, at 1.96ppm), 2.50-2.35 (1H, m), 2.25 -2.10 (1H, m), 1.85-1.70(1H, m).

IR (neat) : 3450, 1640cm<sup>-1</sup>.

HCl salt : mp 195-197.5°C.

5 Anal. Calcd for C<sub>23</sub>H<sub>28</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>2</sub>S HCl 0.5H<sub>2</sub>O : C, 53.86; H, 5.90; N, 5.46.

Found : C, 54.08; H, 5.91; N, 5.39.

### Example 33

(2S,3S)-1-[2-N-(2-N,N-Dimethylaminoethylamino)-2-phenylethyl]-3-metoxymetoxypyrrolidine

10 This compound was prepared in 53.1% yield according to a procedure similar to that described in Example 29.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.39-7.20 (5H, m), 4.65 (1H, d, J=7.0Hz), 4.62 (1H, d, J=6.6Hz), 4.28-4.21 (1H, m), 3.70 (1H, dd, J = 3.7, 10.6Hz), 3.37 (3H, s), 2.95 (1H, dd, J=6.2, 9.9Hz), 2.82 (1H, dd, J=10.6, 11.7Hz), 2.80-2.71 (1H, m), 2.59-2.45 (4H, m), 2.42-

15 2.30 (4H, m), 2.18 (6H, s), 2.16-2.05 (1H, m), 1.85-1.75 (1H, m).

IR (neat) : 3300cm<sup>-1</sup>.

### Example 34

2-(3,4-Dichlorophenyl)-N-(2-N,N-dimethylaminoethyl-amino)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

20 This compound was prepared in 88% yield according to a procedure similar to that described in Examples 1 and 25.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.41-7.08 (8H, m), 6.03 (0.7H, dd, J = 5.9, 9.9Hz), 5.01 (0.3H, dd, J=7.0, 7.3Hz), 4.32-4.22 (1H, m), 3.79 (0.6H, s), 3.77 (0.7H, d, J=15.4Hz), 3.70 (0.7H, d, J = 15.4Hz), 3.36 (0.6H, app.t, J=7.0Hz), 3.20 (1.4H, app.t, J=7.3Hz),  
25 3.15-3.00 (2H, m), 2.95-2.80 (1H, m), 2.75-2.50 (2H, m), 2.50-1.61 (6H, m), 2.11 (1.8H, s), 2.09 (4.2H, s).

IR (neat) : 3400, 1640cm<sup>-1</sup>.

HCl salt : amorphous solid.

Anal. Calcd for C<sub>24</sub>H<sub>31</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub> 2HCl 1.3H<sub>2</sub>O : C, 51.40; H, 6.40; N, 7.49.

30 Found : C, 51.79; H, 7.01; N, 7.58.

### Example 35

(2S,3S)-1-[2-N-(2,2-Dimethoxyethylamino)-2-phenylethyl]-3-hydroxypyrrolidine

This compound was prepared in 66% yield according to a procedure similar to that described in Example 2.

-28-

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.38-7.22 (5H, m), 4.45 (1H, t, J=5.1Hz), 4.33-4.27 (1H, m), 3.74 (1H, dd, J = 3.7, 10.6Hz), 3.36 (3H, s), 3.30 (3H, s), 3.08-3.00 (1H, m), 2.85 (1H, dd, J=10.6, 12.1Hz), 2.71 (1H, br.d, J=9.5Hz), 2.65-2.55 (3H, m, including 2H, d, J=5.1Hz at 2.61ppm), 2.37 (1H, dd, J=3.7, 12.1Hz), 2.32-2.12 (4H, m), 1.81-1.72 (1H, m).

5 IR (neat) : 3400, 3300cm<sup>-1</sup>.

#### Example 36

2-(3,4-Dichlorophenyl)-N-(2,2-dimethoxyethyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

This compound was prepared in 91.2% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.41-7.10 (8H, m), 6.09 (1H, dd, J = 5.9, 9.9Hz), 4.35-4.25 (1H, m), 3.94 (1H, d, J=15.4Hz), 3.85 (1H, d, J=15.8Hz), 3.59 (1H, t, J=5.1Hz), 3.34-3.15 (3H, m), 3.22 (3H, s), 3.18 (3H, s), 3.10-3.00 (1H, m), 2.86 (1H, dd, J=5.9, 12.5Hz), 2.76-2.63 (2H, m), 2.40-2.25 (1H, m), 2.20-2.08 (1H, m), 1.90-1.65 (2H, m).

15 IR (neat) : 3450, 1640cm<sup>-1</sup>.

Fumalic acid salt : amorphous solid.

Anal. Calcd for C<sub>24</sub>H<sub>30</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>4</sub> · C<sub>4</sub>H<sub>4</sub>O<sub>4</sub> · 2H<sub>2</sub>O : C, 53.09; H, 6.05; N, 4.92.

Found : C, 53.47; H, 6.04; N, 4.51.

#### Example 37

20 (2S,3S)-3-Hydroxy-1-[2-phenylethyl-2-N-(2-pyrrolyl)methyl-amino]pyrrolidine

This compound was prepared in 100% yield according to a procedure similar to that described in Example 9.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 9.29 (1H, br.s), 7.37-7.25 (5H, m), 6.75 (1H, d, J=1.5Hz), 6.12 (1H, dd, J=2.6, 5.5Hz), 5.97 (1H, br.s), 4.35-4.25 (1H, m), 3.78 (1H, d, J=13.9Hz), 3.73 (1H, dd, J=3.3, 11.3Hz), 3.55 (1H, d, J=13.9Hz), 3.31 (2H, br.s), 2.92 (1H, dd, J=11.7, 12.1Hz), 2.87-2.77 (2H, m), 2.52 (1H, dd, J=4.8, 9.9Hz), 2.30 (1H, dd, J=3.3, 12.1Hz), 2.25-2.02 (2H, m), 1.83-1.73 (1H, m).

IR (neat) : 3300cm<sup>-1</sup>.

#### Example 38

30 2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(2-pyrrolyl)methylacetamide

This compound was prepared in 81.3% yield according to a procedure similar to that described in Example 1.

-29-

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 12.00 (1H, br.s), 7.45-7.25 (6H, m), 7.08 (1H, d, J=1.5Hz), 6.87 (1H, dd, J=1.5, 8.1Hz), 6.80-6.75 (1H, m), 6.38 (1H, dd, J=3.7, 12.8Hz), 6.13 (1H, dd, J=2.6, 5.1Hz), 5.99 (1H, br.s), 4.58-4.50 (1H, m), 4.15 (2H, s), 3.65-3.40 (4H, m), 3.00-2.90 (2H, m), 2.80-2.65 (2H, m), 2.40-2.25 (2H, m), 1.90-1.65 (2H, m).

5 IR (neat) : 3450, 1630cm<sup>-1</sup>.

Fumalic acid salt : amorphous solid.

Anal. Calcd for C<sub>23</sub>H<sub>27</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub> C<sub>4</sub>H<sub>4</sub>O<sub>4</sub> H<sub>2</sub>O : C, 57.43; H, 5.48; N, 6.93.

Found : C, 57.46; H, 5.41; N, 6.95.

#### Example 39

10 (2S,3S)-3-Hydroxy-1-[2-N-(1-methyl-2-pyrrolyl)methylamino-2-phenylethyl]pyrrolidine

This compound was prepared in 100% yield according to a procedure similar to that described in Example 9.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.43-7.24 (5H, m), 6.55 (1H, dd, J=1.8, 2.6Hz), 6.05-5.99 (2H, m), 4.30-4.25 (1H, m), 3.75 (1H, dd, J=3.7, 10.6Hz), 3.61 (1H, d, J=13.9Hz), 3.55 (3H, s),

15 3.49 (1H, d, J=13.6Hz), 2.87-2.81 (1H, m), 2.81 (1H, dd, J=10.6, 12.1Hz), 2.62-2.53 (2H, m), 2.29 (1H, dd, J=3.7, 12.1Hz), 2.27-2.08 (2H, m), 1.99 (2H, br.s), 1.76-1.65 (1H, m).

IR (neat) : 3250cm<sup>-1</sup>.

#### Example 40

20 2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(1-methyl-2-pyrrolyl)methylacetamide

This compound was prepared in 57.6% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, DMSO-d<sub>6</sub>) δ 7.54 (1H, d, J=8.1Hz), 7.39 (1H, br.s), 7.35-7.20 (5H, m), 7.17 (1H, br.d, J=8.4Hz), 6.59 (1H, br.s), 5.85-5.70 (2H, m), 5.60-5.50 (1H, m), 4.75-4.60

25 (1H, m), 4.42 (2H, ABq), 4.13 (1H, br.s), 3.70 (1H, s), 3.41 (3H, s), 3.10-2.75 (3H, m), 2.65-2.20(4H, m), 2.00-1.85 (1H, m), 1.55-1.45 (1H, m).

IR (neat) : 3450, 1640cm<sup>-1</sup>.

Fumalic acid salt : amorphous solid.

Anal. Calcd for C<sub>23</sub>H<sub>27</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub> C<sub>4</sub>H<sub>4</sub>O<sub>4</sub> 1.5H<sub>2</sub>O : C, 57.24; H, 5.76; N, 6.67.

30 Found : C, 57.17; H, 5.40; N, 6.52.

#### Example 41

(2S,3S)-3-Hydroxy-1-[2-N-(methoxycarbonyl)methylamino-2-phenylethyl]pyrrolidine

This compound was prepared in 81.4% yield according to a procedure similar to that described in Preparation 3.



-30-

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.36-7.24 (5H, m), 4.37-4.30 (1H, m), 3.82 (1H, dd, J=3.7, 11.0Hz), 3.69 (3H, s), 3.39 (1H, d, J=17.9Hz), 3.22-3.15 (1H, m), 3.18 (1H, d, J=17.6Hz), 2.86 (1H, dd, J=11.0, 12.1Hz), 2.73 (1H, dd, 4.8, 9.9Hz), 2.66 (1H, br.d, J=8.1Hz), 2.60-2.15 (5H, m, including 1H, dd, J=3.7, 12.1Hz, at 2.32ppm), 1.82-1.75 (1H, m).

5

**Example 42****2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-methoxycarbonylmethylacetamide**

This compound was prepared in 57.6% yield according to a procedure similar to that described in Example 1.

10 <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.41-7.20 (6H, m), 7.16-7.08 (2H, m), 6.03 (0.5H, dd, J=5.9, 10.3Hz), 4.99 (0.5H, dd, J=6.2, 8.8Hz), 4.28 (0.5H, d, J=16.9Hz), 4.30-4.20 (1H, m), 3.90 (1H, s), 3.87 (0.5H, d, J=16.9Hz), 3.77 (0.5H, d, J=12.5Hz), 3.75 (1H, s), 3.72 (0.5H, d, J=12.5Hz), 3.68 (1.5H, s), 3.63 (1.5H, s), 3.16-2.42 (5H, m), 2.27-1.65 (4H, m).

IR (neat) : 3450, 1750, 1650cm<sup>-1</sup>.

15 HCl salt : mp 169-172°C.

Anal. Calcd for C<sub>22</sub>H<sub>26</sub>Cl<sub>2</sub>N<sub>2</sub>O<sub>4</sub> · HCl 0.5H<sub>2</sub>O : C, 54.08; H, 5.52; N, 5.48.

Found : C, 54.39; H, 5.49; N, 5.53.

**Example 43****(2S,3S)-1-[2-N-(2-Cyanoethylamino)-2-phenylethyl]-3-hydroxypyrrolidine**

20 A mixture of (2S,3S)-1-(2-amino-2-phenylethyl)-3-hydroxypyrrolidine (0.619g, 3mmol) and acrylonitrile (1ml, 15mmol) in ethanol (6ml) was stirred at room temperature for 13h. The solvent was evaporated to give 0.786mg (100%) of a yellow solid.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.39-7.24 (5H, m), 4.37-4.32 (1H, m), 3.76 (1H, dd, J=3.7, 11.0Hz), 3.08-3.00 (1H, m), 2.90-2.72 (3H, m, including 1H, dd, J=11.0, 12.1Hz, at 2.82ppm), 2.69 (2H, d, J=3.7Hz), 2.53-2.14 (7H, m, including 1H, dd, J=3.7, 12.1Hz at 2.34ppm), 1.83-1.73 (1H, m),.

IR (neat) : 3350, 3280, 2250cm<sup>-1</sup>.

**Example 44****N-2-Cyanoethyl-2-(3,4-dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide**

30

This compound was prepared in 93.3% yield according to a procedure similar to that described in Example 29.

-31-

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.42-7.06 (8H, m), 6.11 (0.3H, dd, J=4.8, 11.4Hz), 5.03 (0.7H, dd, J=5.9, 9.2Hz), 4.40-4.28 (1H, m), 3.85-3.40 (4H, m, including 2H, br.s at 3.77ppm), 3.30-2.60 (5H, m), 2.45-2.08(4H, m), 1.90 (1H, br.s), 1.80-1.65 (1H, m).

IR (neat) : 3450, 2250, 1650cm<sup>-1</sup>.

5 Maleic acid salt : amorphous solid.

Anal. Calcd for C<sub>22</sub>H<sub>23</sub>Cl<sub>2</sub>N<sub>3</sub>O<sub>2</sub>·C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>·H<sub>2</sub>O : C, 55.87; H, 5.38; N, 7.24.

Found : C, 56.12; H, 5.35 N, 7.26.

#### Example 45

10 2-(Benzo[b]furan-4-yl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-methoxycarbonylmethylacetamide

This was prepared in 83.7% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>, it appeared as 1:1 rotamer mixture by amide bond) δ 7.64 (0.5H, d, J=2.2Hz), 7.61 (0.5H, d, J=2.2Hz), 7.46-7.40 (1H, m), 7.35-7.20 (6.5H, m), 7.10-7.05 (1.5H, m), 6.88 (0.5H, d, J=2.2Hz), 6.85 (0.5H, d, J=2.2Hz), 6.09 (0.5H, dd, J=5.9, 10.6Hz), 5.09 (0.5H, dd, J=7.3, 8.1Hz), 4.26 (0.5H, d, J=16.5Hz), 4.24-4.10 (1H, m), 4.07-3.84 (3H, m, including 1H, s, at 4.02ppm and 1H, s, at 3.92ppm), 3.81 (0.5H, d, J=16.5Hz), 3.65 (1.5H, s), 3.61 (1.5H, s), 3.18-3.09 (1H, m), 2.95-1.55 (8H, m).

IR (KBr) : 3420, 1740, 1635cm<sup>-1</sup>.

20 Fumalic acid salt : amorphous solid.

Anal. Calcd for C<sub>25</sub>H<sub>28</sub>N<sub>2</sub>O<sub>5</sub>·C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>·0.5H<sub>2</sub>O : C, 62.02; H, 5.92; N, 4.99.

Found : C, 62.08; H, 5.80; N, 4.97.

#### Example 46

25 N-[2-(3-(S)-Hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-methoxycarbonylmethyl-2-(4-trifluoromethylphenyl) acetamide

This was prepared in 87.9% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>, it appeared as 1:1 rotamer mixture by amide bond) δ 7.60 (1H, d, J=8.4Hz), 7.57 (1H, d, J=8.8Hz), 7.43 (1H, d, J=8.1Hz), 7.37 (1H, d, J=8.1Hz), 7.34-7.26 (4H, m), 7.10 (1H, d, J=2.2, 7.7Hz), 6.05 (0.5H, dd, J=5.9, 10.6Hz), 5.00 (0.5H, dd, J=7.0, 8.1Hz), 4.25 (0.5H, d, J=16.8Hz), 4.30-4.15 (1H, m), 3.95-3.73 (3.5H, m), 3.67 (1.5H, s), 3.62 (1.5H, s), 3.20-3.05 (1H, m), 3.00-1.55 (8H, m).

IR (neat) : 3450, 1750, 1650cm<sup>-1</sup>.

Fumalic acid salt : amorphous solid.

-32-

Anal. Calcd for  $C_{24}H_{27}F_3N_2O_4 \cdot C_4H_4O_4 \cdot 0.5H_2O$  : C, 57.04; H, 5.47; N, 4.75.

Found : C, 57.23; H, 5.31; N, 4.70.

**Example 47**

N-[2-(3-(S)-Hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-methoxycarbonylmethyl-2-(3-nitrophenyl)acetamide

This was prepared in 97.8% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>, it appeared as 1:1 rotamer mixture by amide bond)  $\delta$  8.13-8.06 (2H, m), 7.64 (1H, br.d, J=7.0Hz), 7.50 (1H, dd, J=7.7, 8.8Hz), 7.40-7.15 (5H, m), 6.05 (0.5H, dd, J=5.9, 10.6Hz), 5.06 (0.5H, dd, J=6.2, 8.8Hz), 4.29 (0.5H, d, J=16.8Hz), 4.30-4.20 (1H, m), 3.97 (1H, s), 3.97-3.87 (2H, m, including 1H, dd, J=4.0, 9.5Hz at 3.92ppm), 3.78 (0.5H, d, J=15.7Hz), 3.68 (1.5H, s), 3.66 (1.5H, s), 3.20-2.46 (5H, m), 2.32-1.65 (4H, m).

IR (neat) : 3450, 1750, 1650cm<sup>-1</sup>.

Fumalic acid salt : mp 78-80.5°.

Anal. Calcd for  $C_{25}H_{27}N_3O_6 \cdot C_4H_4O_4 \cdot H_2O$  : C, 56.34; H, 5.78; N, 7.30.

Found : C, 56.62; H, 5.83; N, 7.07.

**Example 48**

2-(3-Bromophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-methoxycarbonylmethylacetamide

This was prepared in 82.7% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>, it appeared as 1:1 rotamer mixture by amide bond)  $\delta$  7.48-7.10 (9H, m), 6.05 (0.5H, dd, J=5.9, 10.6Hz), 4.97 (0.5H, dd, J=6.6, 8.4Hz), 4.30 (0.5H, d, J=16.9Hz), 4.27-4.15 (1H, m), 3.89 (1H, s), 3.86 (0.5H, d, J=18.3Hz), 3.80-3.72 (2H, m), 3.69 (1.5H, s), 3.62 (1.5H, s), 3.16-2.52 (4.5H, m), 2.39 (0.5H, dd, J=4.8, 9.5Hz), 2.25-2.00 (2.5H, m), 1.80-1.60 (1.5H, m).

IR (neat) : 3450, 1750, 1650cm<sup>-1</sup>.

Fumalic acid salt : amorphous solid.

Anal. Calcd for  $C_{25}H_{27}BrN_2O_4 \cdot C_4H_4O_4 \cdot H_2O$  : C, 53.21; H, 5.46; N, 4.60.

Found : C, 53.43; H, 5.26; N, 4.36.

**Example 49**

N-[2-(3-(S)-Hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-methoxycarbonylmethyl-2-(2,3,6-trichlorophenyl)acetamide

-33-

This was prepared in 67.2% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>, it appeared as 3:2 rotamer mixture by amide bond) δ 7.45-7.25 (7H, m), 6.04 (0.6H, dd, J=5.9, 10.3Hz), 5.24 (0.4H, dd, J=5.5, 8.8Hz), 4.35-3.80 (5H, m), 3.68 (1.8H, s), 3.63 (1.2H, s), 3.21-3.02 (2H, m), 2.85-2.05 (5H, m), 1.90-1.60 (2H, m).

IR (neat) : 3450, 1740, 1660cm<sup>-1</sup>.

Fumalic acid salt : amorphous solid.

Anal. Calcd for C<sub>23</sub>H<sub>25</sub>CN<sub>2</sub>O<sub>4</sub>·C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>·H<sub>2</sub>O : C, 51.16; H, 4.93; N, 4.42.

Found : C, 51.38; H, 4.96; N, 4.29.

10

#### Example 50

N-[2-(3-(S)-Hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-methoxycarbonylmethyl-2-(1-naphthyl)acetamide

This was prepared in 70.2% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>, it appeared as 1:1 rotamer mixture by amide bond) δ 7.98 (0.5H, d, J=7.7Hz), 7.90-7.75 (2.5H, m), 7.65-7.15 (9H, m), 6.14 (0.5H, dd, J=5.5, 10.6Hz), 5.06 (0.5H, dd, J=6.6, 8.4Hz), 4.40-4.13 (3.5H, m), 3.95 (1H, s), 3.88 (0.5H, d, J=16.9Hz), 3.67 (1.5H, s), 3.62 (1.5H, s), 3.25-1.95 (7.5H, m), 1.80-1.60 (1.5H, m).

IR (KBr) : 3500, 1740, 1630cm<sup>-1</sup>.

20 Fumalic acid salt : mp 189.5-193.5°.

Anal. Calcd for C<sub>27</sub>H<sub>30</sub>N<sub>2</sub>O<sub>4</sub>·C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>·0.5H<sub>2</sub>O : C, 65.14; H, 6.17; N, 4.90.

Found : C, 65.35; H, 6.04; N, 4.91.

#### Example 51

(2S,3S)-1-[2-N-(1,2,4-Oxadiazol-3-yl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine

25 A mixture of 2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethylamine (0.619g, 3mmol), 3-chloromethyl-1,2,4-oxadiazole (533mg, 4.5mmol), and K<sub>2</sub>CO<sub>3</sub> (415g, 3mmol) in DMF (4ml) was stirred at room temperature for 16h. The reaction mixture was poured into water (30ml) and extracted with ethyl acetate (20ml x 3). The extract was washed with brine, dried (Na<sub>2</sub>SO<sub>4</sub>), and concentrated to give 982mg of a yellow oil, which was purified by column chromatography (silica gel; 40g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH: 40/1 to 10/1) to afford 347mg (40.1%) of title compound.

30 <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 8.69 (1H, s), 7.40-7.25 (5H, m), 4.33-4.28 (1H, m), 3.93 (1H, d, J=15.4Hz), 3.79 (1H, d, J=15.4Hz), 3.80-3.75 (1H, m), 3.00-2.86 (2H, m), 2.70-2.60 (2H, m), 2.40-2.31 (3H, m), 2.29-2.10 (2H, m), 1.79-1.70 (1H, m).

#### Example 52

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(1,2,4-oxadiazol-3-yl)methylacetamide

This was prepared in 84.4% yield according to a procedure similar to that described in Example 1.

- 5 <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>, it appeared as 4:1 rotamer mixture by amide bond) δ 8.66 (0.8H, s), 8.60 (0.2H, s), 7.50-7.14 (8H, m), 6.15 (0.8H, dd, J=5.5, 11.0Hz), 5.11 (0.2H, t, J=6.2Hz), 4.90 (0.2H, d, J=16.1Hz), 4.48 (0.2H, d, J=16.1Hz), 4.41 (1.6H, s), 4.25-3.85 (3H, m), 3.27 (0.8H, dd, J=11.7, 12.5Hz), 3.20-1.65 (8.2H, m).

IR (neat) : 3450, 1730, 1650cm<sup>-1</sup>.

- 10 Fumalic acid salt : amorphous solid.

Anal. Calcd for C<sub>22</sub>H<sub>24</sub>C<sub>2</sub>N<sub>4</sub>O<sub>3</sub>·C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>·H<sub>2</sub>O : C, 53.21; H, 4.96; N, 9.19.

Found : C, 53.37; H, 4.87; N, 9.12.

**Example 53**

2-(Benzo[b]furan-4-yl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(1,2,4-oxadiazol-3-yl)methylacetamide

- 15

This was prepared in 61.1% yield according to a procedure similar to that described in Example 1.

- <sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>, it appeared as 3:1 rotamer mixture by amide bond) δ 8.66 (0.75H, s), 8.59 (0.25H, s), 7.62 (1H, d, J=2.2Hz), 7.43 (1H, d, J=8.1Hz), 7.35-7.05 (7H, m), 6.94 (0.75H, d, J=1.1Hz), 6.87 (0.25H, d, J=1.1Hz), 6.19 (0.75H, dd, J=5.5, 11.0Hz), 5.20 (0.25H, d, J=6.2, 8.8Hz), 4.91 (0.25H, d, J=16.1Hz), 4.48 (0.75H, d, J=17.2Hz), 4.41 (0.25H, d, J=16.1Hz), 4.37 (0.75H, d, J=17.2Hz), 4.33 (1.5H, s), 4.20-4.10 (1.5H, m, including 0.5H, s, at 4.11ppm), 3.30 (0.75H, dd, J=11.7, 12.1Hz), 3.20-1.55 (8.25H, m, including 0.75H, dd, J=5.5, 12.5Hz at 2.68ppm).

- 25 IR (neat) : 3450, 1740, 1650cm<sup>-1</sup>.

Fumalic acid salt : amorphous solid.

Anal. Calcd for C<sub>23</sub>H<sub>26</sub>N<sub>4</sub>O<sub>4</sub>·C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>·H<sub>2</sub>O : C, 59.99; H, 5.56; N, 9.65.

Found : C, 59.74; H, 5.26; N, 9.40.

**Example 54**

- 30 (2S,3S)-1-[2-N-(N',N'-Dimethylaminocarbonyl)methylamino-2-phenylethyl]-3-hydroxypyrrolidine

A mixture of (2S,3S)-1-(2-amino-2-phenylethyl)-3-hydroxypyrrolidine (0.413g, 2mmol), 2-chloro-N,N-dimethylacetamide (292mg, 2.4mmol), and K<sub>2</sub>CO<sub>3</sub> (276mg, 2mmol) in DMF (4ml) was stirred at 50° for 2.5h. The reaction mixture was poured into water(10ml) and extracted with ethyl acetate(20ml x 3). After dry(Na<sub>2</sub>SO<sub>4</sub>), the solvent was evaporated to give 558mg of brown

-35-

oil, which was purified by column chromatography (silica gel: 20g, CH<sub>2</sub>Cl<sub>2</sub>/MeOH: 30/1 to 10/1) to give 94.4mg(33.4%) of yellow oil.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.45-7.20 (5H, m), 4.40-4.25 (1H, m), 3.82 (1H, br.d, J=8.4Hz), 3.36 (1H, d, J=16.5Hz), 3.25-3.10 (2H, m, including 1H, d, J=16.1Hz, at 3.17ppm), 3.00-2.05 (17H, m, including 3H, s, at 2.95ppm), 1.90-1.75 (1H, m).  
IR (neat) : 3400, 1640cm<sup>-1</sup>.

**Example 55**

2-(3,4-Dichlorophenyl)-N-(N',N'-dimethylaminocarbonyl)methyl-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]acetamide

10 This was prepared in 64.6% yield according to a procedure similar to that described in Example 1.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>, it appeared as 3:2 rotamer mixture by amide bond) δ 7.41-7.12 (8H, m), 6.09 (0.6H, dd, J=6.4, 9.2Hz), 5.10 (0.4H, t, J=7.3Hz), 4.37 (0.4H, d, J=15.7Hz), 4.30-4.20 (1H, m), 3.97-3.58 (3.6H, m), 3.30-1.70 (15H, m, including 1.2H, s, at 2.98ppm, 1.2H, s, at 2.93ppm, 1.8H, s, at 2.89ppm, 1.8H, s, at 2.86ppm).  
IR (neat) : 3450, 1650cm<sup>-1</sup>.

Fumalic acid salt : amorphous.

Anal. Calcd for C<sub>24</sub>H<sub>29</sub>N<sub>3</sub>O<sub>5</sub>·C<sub>4</sub>H<sub>4</sub>O<sub>4</sub>·1.2H<sub>2</sub>O : C, 54.59; H, 5.79; N, 6.82.

Found : C, 54.81; H, 6.17; N, 6.84.

20

**Example 56**

(2S,3S)-3-Hydroxy-1-[2-N-(6-methylpyridin-2-yl)methylamino-2-phenylethyl]pyrrolidine

This was prepared in 92.8% yield according to a procedure similar to that described in Example 9.

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>) δ 7.49 (1H, t, J=7.7Hz), 7.45-7.25 (5H, m), 7.01 (1H, d, J=7.7Hz), 6.96 (1H, d, J=7.7Hz), 4.35-4.25 (1H, m), 3.81 (1H, d, J=13.9Hz), 3.70 (1H, dd, J=3.7, 11.0Hz), 3.61 (1H, d, J=14.3Hz), 3.05-2.85 (5H, m, including 1H, dd, J=10.6, 12.1Hz), 2.71 (1H, d, J=9.9Hz), 2.65-2.55 (4H, m, including 3H, s at 2.54ppm), 2.33 (1H, dd, J=3.7, 12.1Hz), 2.25-2.10 (2H, m), 1.90-1.70 (1H, m).  
IR (neat) : 3300cm<sup>-1</sup>.

30

**Example 57**

2-(3,4-Dichlorophenyl)-N-[2-(3-(S)-hydroxypyrrolidin-1-yl)-1-(S)-phenylethyl]-N-(6-methylpyridin-2-yl)methylacetamide

This was prepared in 57.6% yield according to a procedure similar to that described in Example 1.

-36-

<sup>1</sup>H NMR (270MHz, CDCl<sub>3</sub>, it appeared as 5:1 rotamer mixture by amide bond) δ 7.45-6.90 (11H, m), 6.20 (0.8H, dd, J=5.1, 9.9Hz), 5.20-5.10 0.2H, m), 4.92 (0.2H, d), 4.43 (1.6H, s), 4.40 (0.2H, d), 4.30-4.23 (0.8H, m), 4.20-4.15 (0.2H, m), 4.02 (0.4H, s), 3.66 (0.8H, d, J=15.4Hz), 3.58 (0.8H, d, J=15.4Hz), 3.30-3.08 (2H, m), 2.80-2.65 (2H, m, including 0.8H, dd, J=5.1, 12.5Hz at 2.75ppm), 2.60-1.55 (8H, m, including 2.4H, s at 2.50ppm).

IR (neat) : 3400, 1650cm<sup>-1</sup>.

HCl salt : amorphous solid.


Anal. Calcd for C<sub>27</sub>H<sub>29</sub>C<sub>2</sub>N<sub>3</sub>O<sub>2</sub>·2HCl·4.4H<sub>2</sub>O : C, 49.84; H, 6.17; N, 6.46.

Found : C, 49.58; H, 6.36; N, 6.86.

-37-

**CLAIMS**

- 5 1. A process of preparing a compound of the formula (I):




and its pharmaceutically acceptable salt, wherein

R is hydrogen or hydroxy;

Ar is phenyl or phenyl substituted with one to three substituents selected from halo, C<sub>1-4</sub> alkyl and C<sub>1-4</sub> alkoxy;

10 X is phenyl or heterocyclic; phenyl or heterocyclic substituted with one to three substituents selected from halo, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy and methoxycarbonyl; mono-, di- or tri-halomethyl; cyano; COR<sup>1</sup>, CH=NOR<sup>2</sup>, OR<sup>2</sup>, SR<sup>2</sup>, CH<sub>2</sub>CN, CH<sub>2</sub>OR<sup>2</sup>, CH<sub>2</sub>SR<sup>2</sup>, CH<sub>2</sub>S(O)R<sup>2</sup>, CH<sub>2</sub>S(O)<sub>2</sub>R<sup>2</sup>, CH<sub>2</sub>(R<sup>2</sup>)R<sup>3</sup>, CH<sub>2</sub>N(R<sup>2</sup>)R<sup>3</sup>, CH<sub>2</sub>NR<sup>2</sup>OH, 15 CH<sub>2</sub>N(COR<sup>2</sup>)OH, CH<sub>2</sub>NR<sup>2</sup>COR<sup>3</sup>, CH<sub>2</sub>NR<sup>2</sup>S(O)<sub>2</sub>R<sup>3</sup> or CH<sub>2</sub>OCOR<sup>2</sup>, wherein R<sup>1</sup> is hydrogen, hydroxy, amino, NHOH, NHOCH<sub>3</sub>, pyridylamino, NHN(CH<sub>3</sub>)<sub>2</sub>, C<sub>1-4</sub> alkoxy, benzyloxy, C<sub>1-4</sub> alkylamino, di-C<sub>1-4</sub> alkylamino, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkylthio; and R<sup>2</sup> and R<sup>3</sup> are each hydrogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy or C<sub>7-11</sub> phenylalkyl; and 20 X<sup>1</sup> is phenyl, naphthyl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl or benzothienyl; phenyl, naphthyl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl or benzothienyl, substituted with one to three substituents selected from halo, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, amino, hydroxy, nitro, trifluoromethyl and mesyl,

which process comprises reacting a compound of the formula (II)






-38-

with an acylating agent in a reaction-inert solvent.

2. A process according to claim 1, the acylating agent is acyl halide.
3. A process according to claim 1, the reaction is carried out at from -30°C to 100°C for 10 minutes to 48 hours.
- 5 4. A process according to claim 1, the solvent is selected from aromatic hydrocarbons, ethers, dioxane and tetrahydrofuran, halogenated hydrocarbons, amides and nitriles.
5. A process according to claim 1, wherein R is hydroxy; Ar is phenyl optionally substituted with one to three halogen atoms; and X<sup>1</sup> is phenyl optionally substituted with one to three halogen atoms.
- 10 6. A process of preparing a compound of the formula (I):




and its pharmaceutically acceptable salt, wherein

- R is hydrogen or hydroxy;
- Ar is phenyl or phenyl substituted with one to three substituents selected from halo, C<sub>1-4</sub> alkyl and C<sub>1-4</sub> alkoxy;
- 15 X is phenyl or heterocyclic; phenyl or heterocyclic substituted with one to three substituents selected from halo, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy and methoxycarbonyl; mono-, di- or tri-halomethyl; cyano; COR<sup>1</sup>, CH=NOR<sup>2</sup>, OR<sup>2</sup>, SR<sup>2</sup>, CH<sub>2</sub>CN, CH<sub>2</sub>OR<sup>2</sup>, CH<sub>2</sub>SR<sup>2</sup>, CH<sub>2</sub>S(O)R<sup>2</sup>, CH<sub>2</sub>S(O)<sub>2</sub>R<sup>2</sup>, CH<sub>2</sub>(R<sup>2</sup>)R<sup>3</sup>, CH<sub>2</sub>N(R<sup>2</sup>)R<sup>3</sup>, CH<sub>2</sub>NR<sup>2</sup>OH,
- 20 CH<sub>2</sub>N(COR<sup>2</sup>)OH, CH<sub>2</sub>NR<sup>2</sup>COR<sup>3</sup>, CH<sub>2</sub>NR<sup>2</sup>S(O)<sub>2</sub>R<sup>3</sup> or CH<sub>2</sub>OCOR<sup>2</sup>, wherein R<sup>1</sup> is hydrogen, hydroxy, amino, NHOH, NHOCH<sub>3</sub>, pyridylamino, NHN(CH<sub>3</sub>)<sub>2</sub>, C<sub>1-4</sub> alkoxy, benzyloxy, C<sub>1-4</sub> alkylamino, di-C<sub>1-4</sub> alkylamino, C<sub>1-4</sub> alkyl or C<sub>1-4</sub> alkylthio; and R<sup>2</sup> and R<sup>3</sup> are each hydrogen, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy or C<sub>7-11</sub> phenylalkyl; and
- X<sup>1</sup> is phenyl, naphthyl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl or benzothienyl;
- 25 phenyl, naphthyl, furyl, thienyl, pyridyl, thiazolyl, benzofuryl or benzothienyl, substituted with one to three substituents selected from halo, C<sub>1-4</sub> alkyl, C<sub>1-4</sub> alkoxy, amino, hydroxy, nitro, trifluoromethyl and mesyl,

which process comprises reacting a compound of the formula (III)

-39-



(III)

with an alkylating agent in a reaction-inert solvent.

7. A process according to claim 6, the alkylating agent is alkylhalide.
8. A process according to claim 6, the reaction is carried out at from 0°C to 200°C for 5  
5 minutes to 24 hours.
9. A process according to claim 6, the solvent is selected from aromatic hydrocarbons, ethers, dioxane and tetrahydrofuran, halogenated hydrocarbons, amides and nitriles.
10. A process according to claim 6, wherein R is hydroxy; Ar is phenyl optionally  
substituted with one to three halogen atoms; and X<sup>1</sup> is phenyl optionally substituted with one to  
10 three halogen atoms.

# INTERNATIONAL SEARCH REPORT

International Application No  
PCT/IB 95/00374

**A. CLASSIFICATION OF SUBJECT MATTER**  
IPC 6 C07D207/12 C07D207/32 C07D401/12 C07D405/12 C07D409/12  
A61K31/40

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)  
IPC 6 C07D

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	EP,A,0 483 580 (MERCK) 6 May 1992 see the whole document ---	1-10
Y	EP,A,0 569 802 (MERCK) 18 November 1993 see the whole document -----	1-10

☐ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \* "A" document defining the general state of the art which is not considered to be of particular relevance
- \* "E" earlier document but published on or after the international filing date
- \* "I" document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \* "(O)" document referring to an oral disclosure, use, exhibition or other means
- \* "P" document published prior to the international filing date but later than the priority date claimed

- \* "T" later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \* "X" document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \* "Y" document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \* "&" document member of the same patent family

Date of the actual completion of the international search

25 September 1995

Date of mailing of the international search report

- 5. 10. 95

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. ( + 31-70) 340-2040, Tx. 31 651 epo nl,  
Fax ( + 31-70) 340-3016

Authorized officer

De Jong, B

**INTERNATIONAL SEARCH REPORT**

Information on patent family members

International Application No.

PCT/IB 95/00374

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP-A-0483580	06-05-92	DE-A- 4034785	07-05-92
		AU-A- 8553891	07-05-92
		CA-A- 2054648	03-05-92
		JP-A- 4264067	18-09-92
		US-A- 5232978	03-08-93
-----			
EP-A-0569802	18-11-93	DE-A- 4215213	11-11-93
		AU-B- 3834193	11-11-93
		CN-A- 1079219	08-12-93
		CZ-A- 9300823	16-02-94
		JP-A- 6049022	22-02-94
		PL-A- 298845	10-01-94
		ZA-A- 9303222	08-12-93
-----			